




BİYOTEKNİKPRODUCT CATALOG





WE WORK FOR ANIMAL HEALTH





medical and non-medical products and premixes since 1955. As separate facilities in Kırklareli Organized Industrial Zone, BTI a domestic and strong brand, Biyoteknik is one of the leading a GMP-certified facility started operations in 2018, and BT2 the veterinary products manufacturers in Türkiye, standing out with second facility for the production of non-medical products is its know-how in the sector and increasing its market share every commissioned in 2021. year with its broad product range.

network in national and international markets

Biyoteknik has a long-established history in the field of veterinary Biyoteknik continues its production and R&D activities with two

1955'ten

The vision of the company is to ensure sustainability by protecting The company serves its clients with a wide sales and distribution natural resources and the ecosystem and to carry the brand to future generations.



Our Quality and Service Standards

Biyoteknik holds ISO 9001-2015 and ISO 22000-2018 quality management system certificates.

Our corporation holds the "Good Manufacturing Practices" (GMP) certificate issued by the Ministry of Agriculture and Forestry. We manufacture veterinary pharmaceuticals according to GMP standards.

ucts, feed additives/premixes and veterinary pharmaceuticals are approved by the Min-

Biocidal products are manufactured and pre- private label clients feel comfortable to rely on sented for sale with the approval of Ministry of our products and manufacturing expertise for Health. Biyoteknik manufactures each one of private label manufacturing partnership. its products to the highest quality standards Non-medicinal veterinary health prod- by applying the quality assurance system effectively across all of its processes.

Private Label

We have broad and in-depth expertise in private label manufacturing. We understand how to address our customers needs for customization, speed to market, quality assurance and cost pressure. Our wide range of products have a istry of Agriculture and Forestry whereas the proven track record in the market. Therefore our

Contract Manufacturing

Our veterinary pharma manufacturing experience dates back to 1955. Since then, we have developed compelling manufacturing skills and infrastructure to offer seamless manufacturing services. We offer to our contract manufacturing clients state of the art facilities, flexible manufacturing solutions, cost-effective manufacturing expertise and dedication to quality assurance. Building partnership with us for your contract manufacturing needs means gaining a manufacturing partner who dedicates his resources to your success.











Wide Sales Network

We have a wide geographic sales network and team are great assets supporting our strong market logistics efficiency, CRM applications, warehousing our sales structure. and well-established client relationship of our sales

coverage in Türkiye for veterinary pharmaceuticals. If position at sales in Türkiye. Our active client database you plan to expand your geographic sales network in and our long term client relationship with both Türkiye, we offer great partnership opportunity. Our distributors and veterinary clinics are backbone of









Quality Control

The quality control division consists of Physico-Chemicial Analysis Laboratory, Microbiology Analysis Laboratory and Research and Development Laboratory.

With our experienced team we perform material as per specifications. quality control analyses according to preset

specifications for all constituents such as raw material and packaging procured for production as part of quality management system. Manufacturing process starts with approved raw







R&D and Innovation

new products perusing the scientific and relevant regulations. statistical data, conducting literature review, and monitoring technological advancements in the great emphasis on R&D works. The R&D sector. We continuously enhance the existing team improves the product range day by products within our portfolio. We formulate new day, by adding new products developed and innovative products to satisfy the need of the according to the latest trends at home and sector and decrease the dependency on imported abroad. goods. Quality control, R&D and Regulatory Affairs departments collaborate with each other to

The team of specialists at R&D develops ensure the compliance of our products with

Biyoteknik steers the sector by placing







| VETERINARY MEDICINAL PRODUCTS | | BİYO DERMACURE CURLY SKIN CARE PRODUCT FOR CATS AND DOGS CURLY DOGS | 34 |
|---|----|---|------|
| BİYOTOKS CONCENTRATE FOR DIP SOLUTION - VETERINARY ECTOPARASITICIDE (Insecticide - Acaricide) | 11 | BİYO DERMACURE FRESH DEODORANT SHAMPOO SKIN CARE PRODUCT FOR DOG | 35 |
| DEKATOKS CONCENTRATE FOR DIP EMULSION - VETERINARY ECTOPARASITICIDE (Insecticide - Acaricide | | BIYO DERMACURE WHITE SKIN CARE SHAMPOO FOR WHITE DOGS | 36 |
| FLUMET POUR-ON SOLUTION - ECTOPARASITICIDE (Insecticide - Acaricide) | 13 | BİYO DERMACURE BLACK SKIN CARE SHAMPOO FOR BLACK DOGS | 37 |
| | | BİYO DERMACURE COMFORT SKIN CARE SHAMPOO FOR CATS & DOGS | 38 |
| BİYOLİNE KEDİ SPOT-ON SOLUTION ECTOPARASITIC (Insecticide and Acaricide) for Cats | 14 | BİYO DERMACURE PUPPY SKIN CARE SHAMPOO FOR PUPPIES | 39 |
| BİYOLİNE KÖPEK SPOT-ON SOLUTION ECTOPARASITIC (Insecticide and Acaricide) for Dogs | 15 | BİYO DERMACURE PET-DUST DRY POWDER DRY SHAMPOO SKIN CARE PRODUCT FOR CATS AND DOG | S 40 |
| İMİDA SPOT-ON SOLUTION ECTOPARASITIC (Insecticide) | 16 | | |
| IMIDAMIX SPOT-ON SOLUTION ECTOPARASITIC (Insecticide) for Dogs | 17 | BİYO DERMACURE HERBİO DRY FOAM DRY FOAM SHAMPOO FOR DOGS AND CATS | 41 |
| BIYOCIDAL PRODUCTS | | BİYO DERMACURE DRY FOAM SKIN CARE PRODUCT FOR CATS AND DOGS | 42 |
| BITOCIDAL PRODUCTS | | SKIN CARE | |
| SİPERTOKS EMULSION CONCENTRATE (EC) | 19 | BİYO DERMACURE CAT-LİNE SOLUTION | 44 |
| PERGRAN GRANULE | 20 | BIYO DERMACURE DOG-LINE SOLUTION | 45 |
| BİYOPET AEROSOL | 21 | BİYO DERMACURE HERBİO CATSPRAY PUMP LIQUID SPRAY® | 46 |
| WELLNESS & CROOMING | | BİYO DERMACURE HERBİO DOGSPRAY PUMP LIQUID SPRAY ® | 47 |
| WELLNESS & GROOMING | | VERABLU AEROSOL SPRAY - SKIN CARE PRODUCT ® | 48 |
| POMADES | | KREOLÍN NAIL CARE SOLUTION | 49 |
| BİYO ICHTHYODERM SKIN CARE POMADE | 23 | BORDUST SKIN CARE PRODUCT | 50 |
| BİYO İYODERM SKIN CARE POMADE | 24 | EYE & EAR CARE | |
| BİYO KAMFODERM SKIN CARE POMADE | 25 | BİYO DERMACURE EAR CLEANER EAR CARE SOLUTION - FOR CATS AND DOGS | 52 |
| BİYO SALİDERM SKIN CARE POMADE | 26 | BİYO DERMACURE EYE CLEANER EYE CARE SOLUTION - FOR CATS AND DOGS | 53 |
| BİYO RİVODERM SKIN CARE POMADE | 27 | ORAL CARE | |
| BİYO ZİNCODERM SKIN CARE POMADE | 28 | VERAXIN SPRAY SOLUTION - ORAL AND DENTAL CARE PRODUCT | 55 |
| BİYO DERMACURE TRİCOBOR SKIN CARE POMADE | 29 | HYGIENE PRODUCT | |
| VERAJEL SKIN CARE PRODUCT | 30 | BİYO TRAİNER SPRAY FOR TOILET TRAINING OF CATS AND DOGS | 57 |
| SHAMPOOS | | SUPPLEMENTS FOR PETS | |
| BIYO DERMACURE TRICOBOR SHAMPOO | 32 | BİYO POWERCURE DOG-VİT ONE-A-DAY MULTIVITAMIN TABLET FOR DOGS ® | 59 |
| BIYO DERMACURE HERBIO DRY FOAM SHAMPOO FOR DOGS AND CATS | 33 | BİYO POWERCURE CAT-VİT ONE-A-DAY MULTIVITAMIN TABLET FOR CATS ® | 60 |

| BIYO POWERCURE CALCIVIT-D ONE-A-DAY CALCIUM-PHOSPHORUS-D3 FOR CATS AND DOGS3 | 61 |
|---|----|
| BÍYO POWERCURE BREWERS YEAST ONE-A-DAY PREMIX GARLIC YEAST TABLET - FOR CAT AND DOG | 62 |
| BİYO POWERCURE METHİO ONE-A-DAY AMINO ACID AND VITAMIN TABLET FOR CATS AND DOGS | 63 |
| BIYO POWERCURE BIYOFLEX ONE-A-DAY JOINT HEALTH SUPPLEMENT FOR CATS AND DOGS ® | 64 |
| BIYO POWERCURE BIYOTAN ONE-A-DAY VITAMIN TABLET FOR CATS AND DOGS ® | 65 |
| BÍYO POWERCURE CAT-FORMULA COMPLEMENTARY FEED FOR KITTENS | 66 |
| BIYO POWERCURE DOG-FORMULA COMPLEMENTARY FEED FOR PUPPIES | 67 |
| BİYO-ANİSOL PET VITAMIN - AMINO ACID | 68 |
| ANTI HAIRBALL PRODUCTS | |
| FERMACAT ANTI - HAIRBALL CAT MALT | 70 |
| FELAXA BİYO-ZINC ANTI-HAIRBALL AND ZINC-CONTAINING MALT FOR CATS | 71 |
| FELAXA JOINT JOINT - BONE-MUSCLE SUPPORTING MALT | 72 |
| FELAXA MULTIVITAMIN MULTIVITAMIN + AMINOACID MALT | 73 |
| FELAXA TAURINE ANTI-HAIRBALL AND TAURINE-CONTAINING MALT | 74 |
| FELAXA LYSINE ANTI - HAIRBALL AND LYSINE - CONTAINING MALT FOR CATS ® | 75 |
| SUPPLEMENTS FOR FARM ANIMALS | |
| VIOMIX VITAMIN-MINERAL FOR RUMINANTS | 77 |
| MINEMIX GOLD VITAMIN - MINERAL - PROBIOTIC FOR RUMINANTS | 78 |
| MİNEMİX VITAMIN - MINERAL - PROBIOTIC FOR RUMINANTS | 79 |
| BİYO ANİSOL SOLUTION VITAMIN - AMINO ACID | 80 |
| KALFOMÍN SOLUTION LIQUID AMINO ACID AND MINERAL | 81 |
| BİYOMİX COMPLEMENTARY FEED FOR RUMINANT | 82 |
| BİYOSACC VITAMIN, MINERAL, PROBIOTIC PREMIX | 83 |
| VIOMIN-B COMPLEMENTARY FEED FOR RUMINANT | 84 |
| VÍOMÍN-E VITAMIN, MINERAL, AMINOACID PREMIX FOR RUMINANTS | 85 |

| VİTHORSE VITAMIN - MINERAL - AMINOACID | 86 |
|--|-----|
| BİYO EVİT-SE VITAMIN - MINERAL COMPLEMENTARY FEED | 87 |
| BİYO AVİT-K VITAMIN COMPLEMENTARY FEED | 88 |
| VÍOSOL-K MULTIVITAMIN POWDER FOR POULTRY | 89 |
| PRODIGEST INTESTINAL FLORA REGULATOR PROBIOTIC AND MINERAL PREMIX | 90 |
| BİYO SEL-DE CARLSBAD POWDER COMPLEMENTARY FEED | 91 |
| BİYO MAGNESİE CALCINEI POWDER COMPLEMENTARY FEED | 92 |
| BİYO MAGNEZYUM SÜLFAT POWDER COMPLEMENTARY FEED | 93 |
| BİYOCAL SOLUTION COMPLEMENTARY LIQUID MINERAL FEED | 94 |
| BISMOLIN ORAL POWDER - COMPLEMENTARY FEED | 95 |
| ARISOL MULTIVITAMIN FOR HONEYBEES | 96 |
| ORNAMENTAL AND CAGE BIRDS | |
| BİTTER BEAK ANTI CANNIBAL SPRAY | 98 |
| AVISENCID SKIN CARE SOLUTION | 99 |
| BİYO DERMACURE HERBİO AVİSPRAY PUMP LIQUID SPRAY SKIN CARE CREAM | 100 |
| AVI-AST SOLUTION | 10 |
| AVİ-DİAR SOLUTION PREMIX | 102 |
| AVİSOL-A VITAMIN A - VITAMIN D ₃ - VITAMIN E - VITAMIN C SOLUTION | 103 |
| AVİSOL-B VITAMIN SOLUTION FOR BIRDS | 104 |
| AVİSOL-PLUS VITAMIN SOLUTION FOR BIRDS | 105 |
| AVİSOL-SE MINERAL SOLUTION FOR BIRDS | 106 |
| AVIMINERAL MINERAL SOLUTION FOR BIRDS | 107 |
| AVİ-VİT MULTIVITAMIN FOR BIRDS | 108 |
| FISH AND AQUARIUM | |
| BİYO-AQUACURE CONTRABLUE SOLUTION - AQUARIUM WATER REGULATOR | 110 |
| BİYO-AQUACURE CONTRA SOLUTION - AQUARIUM WATER REGULATOR | 111 |

BİYO-AQUACURE NİTROFUR SOLUTION - AQUARIUM WATER REGULATOR





CONCENTRATE FOR DIP SOLUTION

VETERINARY ECTOPARASITIC (Insecticide - Acaricide)

100 ml - 500 ml - 1000 ml - 5 kg

COMPOSITION: 100 ml of Biyotoks, with a light yellow color and unique odor, contains 7.5 g Cypermethrin (cis:trans 50:50). **INDICATIONS FOR USE:** It is used against scabies, ticks and blood sucking lices in the treatment of the following ectoparasites in cattle, sheep, goats, horses and dogs.

| Ticks (Ixodidae) | | | Scabies | Lices |
|------------------|--------------------|-------------------|-----------------|-------------------|
| Amblyomma spp. | Haemaphysalis spp. | Ixodes spp. | Psoroptes spp. | Haematopinus sp |
| A.Variegatum | H.Otophila | I.Ricinus | Sarcoptes spp. | Linognathus spp. |
| | H.Sulcata | I.Hexagonus | Chorioptes spp. | L. Ovillus |
| Boophilus spp. | H.Numidiana | - | | L. Vituli |
| B.Annulatus | | Rhipicephalus | In dogs: | |
| | Hyalomma spp. | R.Sanguineus | Demodex canis | Damalinia spp. |
| Dermacentor spp. | H.Anatolicum | R.Turanicus | | D. Bovis |
| D.Marginatus | H.Detritum | | | D. Ovis |
| - | | Ornithodoros spp. | | Melophagus Ovinus |
| | | O.Lahorensis | | . 9 |

USAGE AND DOSAGE: Biyotoks is applied externally in the form of bathing, spraying and washing. Biyotoks is used by mixing with water at the rates shown in the table below to obtain a homogeneous mixture. Unless recommended otherwise by the veterinarian, the dose is as indicated in the table.

| Animal Species | Parasite Species | Method of Administration | Biyotoks Administration |
|------------------------------------|--|--|--|
| Cattle, Sheep, Goat Horse | Lice, Flea, Tick, Scabies, Face Fly, Horn Fly, Horse Fly, Ear Tick, Barn Fly, Meat Fly, Mosquito | 3 | 20 ml for 20 liters of water 200 ml for 200 liters of water 1000 ml for 1000 liters of water |
| | Horn Fly, Face Fly, Ear Tickle | 15 ml per ear, 60-120 ml on the face, 360-480 ml along the back are dministered locally with a low-pressure spray. | 75 ml for 5 liters of water |
| Dogs | Ticks, Scabies, Lice, Fleas and all species of flies | The animal is completely wetted by spraying, bathing or wiping with a sponge. It is left to dry by itself in a warm place without rinsing and drying with a towel. Thus, re-infestation is prevented. Repeat after 2-3 weeks if necessary. | |

WITHDRAWAL PERIODS: During the treatment and after the last drug administration, cattle, sheep and goats raised for meat should not be sent to slaughter before 14 days. During the treatment and after the last drug administration, milk obtained before 15 days (30 milkings) in dairy cows and 10 days (20 milkings) in sheep and goats should not be offered for human consumption. Due to the long drug residue purification period in milk; It is not recommended to be applied to cows, sheep and goats producing milk for human consumption.

USE DURING PREGNANCY: It is not used in pregnant animals less than 28 days after birth.

PACKAGING: 100 ml, 500 ml, 1000 ml, 5 kg SHELF LIFE: 5 years







DEKATOKS











CONCENTRATE FOR DIP SOLUTION VETERINARY ECTOPARASITIC

(Insecticide-Acaricide) 100 ml - 500 ml - 1000 ml - 5 lt COMPOSITION: Concentrate for clear, yellow-brown colored immersion emulsion. DEKATOKS contains 50 mg deltamethrin per ml.

INDICATIONS FOR USE: In cattle and sheep; It is used in the fight against ticks, nymphs, larvae, lice, scabies, tsetse fly and common uses, and infestation with fly larvae.

USAGE AND DOSAGE: DEKATOKS concentrate is applied by dipping, spraying or washing. Unless recommended otherwise by the veterinarian, the doses in the table below are applied.

| Cattle | 0.25-1.5 mg/kg body weight can be repeated every 3-6 weeks if necessary. |
|--------|---|
| Sheep | 0.94-4.5 mg/kg body weight can be repeated after 10-21 days if necessary. |

Preparation of Dip Solution; 1 liter of DEKATOKS is used for every 1000 liters of water in the dipping pool. Preparation of Washing and Spraying Solution; 1 ml DEKATOKS is used per 1 liter of water.

WARNINGS ABOUT DRUG RESIDUES IN FOOD: Drug Residue Elimination Time (I.K.A.S.); During the treatment and after the last drug administration, cattle should not be sent to slaughter before 3 days and sheep 7 days. Cow's milk obtained during and after drug use can be offered for human consumption. It should not be used in dairy sheep whose milk is offered for human consumption.

PACKAGING: 100 ml, 500 ml, 1000 ml, 5 lt SHELF LIFE: 3 years



FLUMET











POUR-ON SOLUTION ECTOPARASITIC (Insecticide-Acaricide) 100 ml - 250 ml - 500 ml - 1000 ml **COMPOSITION:** 1 ml of FLUMET Pour-on Solution contains 10 mg of Flumethrin. It is a yellowish light brown solution.

INDICATIONS FOR USE: FLUMET Pour-on Solution is used for ticks, lice and scabies in cattle and sheep during their mature and developmental stages.

| Ticks | Sucking and Biting Lice | Scabies | |
|------------------------|-------------------------|----------------|--|
| Boophilus spp. | Haematophinus sternus | Psoroptes ovis | |
| Hyalomma spp. | Linognathus vituli | Demodex canis | |
| Rhipicephalus spp. | Bavicola bovis, | | |
| Amblyomma spp. | B. ovis, | | |
| Dermacentor marginatus | Melophagus ovinus | | |

USAGE AND DOSAGE: The application dose of FLUMET Pour-on Solution varies according to the type of parasite in the animal. Unless the veterinarian recommends otherwise, a dose of 1 mg/kg body weight against ticks and stinging lice, and 2 mg/kg body weight against sucking lice and psoroptes mange is given.

In cattle: FLUMET Pour-on Solution is poured on the animal's spine from the shoulder to the coccyx. The doses indicated below can be applied in the correct scales with the help of the measuring instrument on the medicine box.

| Cattle Body Weight | FLUMET |
|--------------------|--------|
| Less than 100 kg | 10 ml |
| 100-200 kg | 20 ml |
| 200-300 kg | 30 ml |
| 300-400 kg | 40 ml |
| More than 400 kg | 50 ml |



In sheep: There are two ways to use FLUMET Pour-on Solution in sheep.

1. It is poured to newly sheared sheep on the backbone, from the shoulder to the tailbone, as in cattle. If the fleece or hair of the sheep is long, the fleece or hair on the back is opened by hand and the medicine is poured on the skin part opened in this line.

2. Another way of using it is by placing the animal on its hip. When the animal is seated on its buttocks, the drug is poured into as many points as possible on the fleece or hairless areas on the sides of the abdomen and between the front and hind legs. For such use, a 5 ml plastic syringe without a needle is sufficient. The dose to be administered is withdrawn with a syringe and administered to the animal as described above.

| Sheep Body Weight | FLUMET |
|-------------------|--------|
| 10 kg | 1.0 ml |
| 15 kg | 1.5 ml |
| 20 kg | 2.0 ml |
| 30 kg | 3.0 ml |
| 40 kg | 4.0 ml |
| 50 kg | 5.0 ml |

In Psoroptic Scabies: If there is this type of scabies and lice in cattle and sheep, twice the above-mentioned doses are applied. In dogs: 1 ml/10kg body weight dose is applied in the fight against Demodex scabies. It can be applied directly to the scaly areas or poured on the spine from the back of the head to the coccyx.

USE IN PREGNANCY: No complications were encountered for use during pregnancy.

WARNINGS ABOUT DRUG RESIDUES IN FOOD: Drug Residue Elimination Time (i.k.a.s): Residue wash-out period for meat and milk in cattle and meat in sheep is zero (0). Not applicable to sheep producing milk for human consumption.

PACKAGING: 100 ml, 250 ml, 500 ml, 1000 ml SHELF LIFE: 2 years.

BİYOLİNE KEDİ













SPOT-ON SOLUTION ECTOPARASITIC
(Insecticide and Acaricide) for Cats
0,50 ml

COMPOSITION: It contains 10 g fipronil and 12 g s-methoprene per 100 ml.

INDICATIONS FOR USE: It is used against external parasites in cats, which are sensitive to Fipronil and (S)-Methoprene, as indicated in the table below.


| Flea | Tick | Biting Lice |
|-----------------------|--|----------------------|
| Ctenocephalides felis | lxodes ricinus Rhipicephalus sanguineus | Felicola subrostatus |

It is used in the prevention and treatment of flea, tick infestations, and the destruction of biting lice. In addition, those who come into contact with the drug during the 6 weeks following the application: In stopping the embryonal development (ovicidal effect) in the eggs of adult fleas, It is used to control flea allergic dermatitis (PAD).

USAGE AND DOSAGE: It is applied topically directly on the skin. The hairs between the shoulders of the upright cat are opened so that the skin can be seen. The mouth of the tube is emptied between the shoulders in 2 equal amounts or in lines, in contact with the skin. The minimum therapeutic dose administered pharmacologically in cats is 5 mg/kg body weight for Fipronil and 6 mg/kg body weight for (S)-Methopiene. Practical application doses are adjusted to be above this level. regardless of their body weight it is 1 micro tube of 0.5 ml is for all cats.

USE IN PREGNANCY:Laboratory studies have shown that Fipronil and (S)-Methoprene do not have any teratogenic and embryotoxic effects. When administered multiple times in succession to pregnant and lactating cats at 3 times the recommended dose, the animals did not show any symptoms. This demonstrates the safety of the drug. Biyoline Kedi Spot-on Solution is safely applied to female cats during mating, pregnancy and lactation periods.

PACKAGING: 0.50 ml SHELF LIFE: 5 years



BİYOLİNE KÖPEK











SPOT-ON SOLUTION ECTOPARASITIC (Insecticide and Acaricide) for dogs 0,67 ml - 1,34 ml - 2,68 ml - 4,02 ml COMPOSITION: It contains 10 g fipronil and 9 g s-methoprene per 100 ml.

| Flea | Tick | Biting Lice |
|-----------------------|--|--------------------|
| Ctenocephalides canis | Ixodes ricinus Rhipicephalus sanguineus | Trichodectes canis |
| | Dermacentor marginatus | |

INDICATIONS FOR USE: It is used against external parasites in dogs, which are sensitive to Fipronil and (S)-Methoprene, as indicated in the table below. It is used in the prevention and treatment of flea, tick infestations, and the destruction of biting lice. In addition, those who come into contact with the drug during the 8 weeks following the application: In stopping the embryonal development (ovicidal effect) in the eggs of adult fleas, In stopping the development (larvicidal effect) in the larval and pupal stages, It is used to control flea allergic dermatitis (PAD).

USAGE AND DOSAGE: It is applied topically directly on the skin. The hair between the shoulders of an upright dog is opened so that the skin can be seen. The mouth of the tube is emptied equally in 4 different places from the shoulder to the coccyx, in contact with the skin. The minimum pharmacologically administered therapeutic dose in dogs is 6.7 mg/kg bw for Fipronil, 6 mg/kg bw for (S)-Methoprene. Practical application doses are 0.67 ml, 1.34 ml and 2.68 ml according to the weight of the dog.

| Animal Weight | Applied Dose Fipronil | (S) Methyloprene | Microtube |
|-----------------|-----------------------|-----------------------|-----------|
| Less than 10 kg | 67 mg Fipronil | 60 mg (S)-Methoprene | 0,67 ml |
| 0 - 20 kg | 134 mg Fipronil | 121 mg (S)-Methoprene | 1,34 ml |
| 20- 40 kg | 268 mg Fipronil | 240 mg (S)-Methoprene | 2,68 ml |
| 40- 60 kg | 402 mg Fipronil | 360 mg (S)-Methoprene | 4,02 ml |

USE IN PREGNANCY: Laboratory studies have shown that Fipronil and (S)-Methoprene do not have any teratogenic and embryotoxic effects. When given multiple times in succession at 3 times the recommended dose to pregnant and lactating dogs, the animals showed no symptoms. This demonstrates the safety of the drug. BIOLINE Köpek Spot-on Solution is safely applied to female dogs during mating, pregnancy and lactation periods.

PACKAGING: 0,67 ml - 1,34 ml - 2,68 ml - 4,02 ml


SHELF LIFE: 5 years



IMIDA















SPOT-ON SOLUTION ECTOPARASITIC (Insecticide) for Cats and Dogs 0,4 ml - 1 ml - 2,5 ml - 4 ml - 30 ml COMPOSITION: 100 ml contains 10 g Imidacloprid. It is a colorless to very light yellowish, oily homogeneous solution.

INDICATIONS FOR USE: Imida Spot-on Solution destroys flea infestations of dogs and cats in single application and prevents new flea infestations for at least 4 weeks. The fleas that live on the animal die completely one day after the drug application and a protection period of four weeks is provided.

USAGE AND DOSAGE: Imida Spot-on Solution is in the form of a micro tube and is very easy to apply. The hair between the shoulders of the uprighted cat or dog is opened so that the skin can be seen. Micro tube is discharged between the shoulders in equal amounts as a line or at 2 different points, in contact with the skin. Dispose only on healthy skin. The dose corresponding to the body weight of the animals are as below:

| Body Weight/Animal | Micro-tube | Dropping Bottle | mg/kg Body Weight |
|--------------------|-----------------------|-----------------|-------------------|
| 0-4 kg/cat and dog | 1 x 0.4 ml micro tube | 8 drops | > 10 mg |
| > 4 kg/cat | 2 x 0.4 ml micro tube | 16 drops | < 20 mg |
| 4-10 kg/dog | 1 x 1.0 ml micro tube | 20 drops | 10-25 mg |
| 10-25 kg/dog | 1 x 2.5 ml micro tube | 50 drops | 10-25 mg |
| 25-40 kg/dog | 1 x 4.0 ml micro tube | 80 drops | 10-16 mg |
| > 40 kg/dog | 2 x 4.0 ml micro tube | 160 drops | < 20 mg |

Imida Spot-on Solution destroys fleas on dogs and cats within 24 hours after administration. A single administration prevents new flea infestations for at least 4 weeks. For the most ideal protection and fight against fleas, Imida Spot-on Solution should be applied once a month. Imida Spot-on Solution maintains its effectiveness even if the animal is bathed, exposed to rain or sunlight. If a repeated dose is required before 4 weeks, it should not be administered more than once a week.

USE IN PREGNANCY: It is safe to use on pregnant dogs and cats. **PACKAGING:** 0.4 ml, 1 ml, 2.5 ml, 4 ml , 30 ml



IMIDAMIX















SPOT-ON SOLUTION ECTOPARASITIC (Insecticide-Acaricide) for Dogs 1 ml - 2,5 ml - 4 ml - 30 ml **COMPOSITION:** 1 ml of IMIDAMIX Spot-on Solution contains 100 mg of Imidacloprid and 500 mg of Permethrin. It is a white, oily solution with an aromatic odour.

INDICATIONS FOR USE: IMIDAMIX Spot-on Solution is used against external parasites such as ticks, fleas, mosquitoes and simuliums in dogs. After application to the skin of dogs, the solution quickly disperses over the body surface. Both active substances are effective on the skin and hair of the subject animal for 4 weeks. External parasites effected by the drug are shown below.

| Parasite Type | e | indication |
|---------------|---|---|
| Ticks | Ixodes rhicinus Rhipicephalus sanguineus Dermacentor spp. | It is used for treatment and protection against ticks. When applied with ticks on dogs, it kills ticks, but these ticks may remain attached to the skin. It also has a repellent and lethal effect against ticks that will jump on animals. It provides efficacy against ticks for 4 weeks with a single treatment. |
| Fleas | Ctenocephalides spp | It is effective against flea infestations for 4 weeks and kills all existing fleas within 12 hours after application. |
| Flies | Mosquito Pheiotomus papatasi (simulium, sand fly) | It has a repellent effect against these fly species. Provides effectiveness for 4 weeks with a single application. |

Indication

USAGE AND DOSAGE: The hair between the shoulders of the animal standing upright is divided into two sides so that the skin is visible. The tip of the micro tube is held to the skin and the micro tube is discharged into the skin equally from 4 different places from the shoulder to the coccyx. Only pour into the healthy part of skin.

Unless recommended otherwise by the veterinarian, the dose is as indicated in the table below.

| Dogs Body Weight | Dose to be a Volume | dministered Drops | Imidacloprid mg/kg body weight | Permethrin mg/kg body weight |
|------------------|---------------------|----------------------|-----------------------------------|---------------------------------|
| 4 kg - 10 kg | 1,0 ml | 20 drops | 10-25 | 50-125 |
| 10 kg - 25 kg | 2,5 ml | 50 drops | 10-25 | 50-125 |
| 25 kg - 40 kg | 4,0 ml | 80 drops | 10-16 | 50-80 |

PACKAGING: 1 ml, 2.5 ml, 4 ml, 30 ml

SHELF LIFE: 5 years

Daracite Type







FORMULATION FORM: EMULSION CONCENTRATE (EC)

100 ml - 500 ml - 1000 ml - 5 lt

COMPOSITION:

Name and amount of active substance (%) (Chemical name; CAS Number)

Cypermethrin cis:trans 50:50 (Cas No: 52315-07-8) 7.5% w/w

Excipients and their amounts (%)

Solvent 88.5% w/w

Emulsifier 4% w/w

| Pest type- life stage | Application area and form | Administration dosage | Range of application |
|-----------------------|--|--|----------------------|
| Tick | Spraying or pulverizing in empty | 26.6 ml/m² of product is | When needed |
| Lice | closed barns, pens, animal shelters, poultry houses and perches | applied from a 1% solution (1L of product is added to 99L | |
| Flea | | of water). (Cypermethrin/m²) | |

Sipertoks is applied in animal shelters such as barns, pens, boxing, poultry coops and perches, in the form of direct spraying for ticks and fleas, or by pulverization. It is suitable for use by adding water at the rate specified in the table according to the application area. During the application, animals, feeders and water bowls should not be kept in the shelters and poultry houses. After being kept closed for 30 minutes, it is ventilated for 1 hour.

PACKAGING: 100 ml, 500 ml, 1000 ml, 5 Lt

SHELF LIFE: 30 months





PERGRAN GRANULE



PERGRAN GRANÜLE 100 gr - 250 gr - 500 gr - 1000 gr

COMPOSITION:

Name and amount of active substance (%) (Chemical name; CAS and EC Number)

Permethrin cis:trans 40:60 (Cas No: 52645-53-1) 0.50 % a/a

Excipients and their amounts (%)

Filler 99.50 % a/a

Pergran is applied as direct sprinkling in animal shelters such as barns, corrals, boxing, poultry houses and roosts for the control of the creatures specified in the table. It is suitable to use in the amounts specified in the table according to the application area. Feeders and drinkers should not be kept in shelters and poultry houses during application. After the application, ventilate the empty closed area for half an hour after being closed for 30 minutes. And clean the product left over from the application. Prevent animals from contacting or eating the product.


| Pest Type-Life stage | Application area | Dosage | Active material dose | Application range |
|----------------------|------------------------------|-------------------------------|----------------------|-------------------|
| Tick | It is applied by sprinkling. | 20 g product / m² | 0,1 g a.i./m² | When necessary |
| Lice | | 20 g product / m ² | 0,1 g a.i./m² | |
| Flea | | 20 g product / m ² | 0,1 g a.i./m² | |
| Ant | | 30 g product / m² | 0,15 g a.i./m² | |

PACKAGING: 100 g - 250 g - 500 g - 1000 g

SHELF LIFE: 2 years



BİYOPET AEROSOL



BİYOPET AEROSOL 150 ml

COMPOSITION:

Name and amount of active substance (%) (Chemical name; CAS Number)

Tetramethrin (Cas No: 7696-12-0) % 0,2 a/a

Piperonyl Butoxide (Cas No: 51-03-06) %2 a/a

Excipients and their amounts (%)

Solvent % 37,8 a/a

Propellant % 60 a/a

Biyopet Aerosol is used in cat, dog and pigeon shelters, empty indoor applications. Care should be taken that there are no animals in the environment. After the application, it is appropriate to keep the application area closed for 30 minutes and to open it after 1 hour of ventilation. During the application, food and water containers should not be kept inside the shelters.

| Pest type- life stage | Application area and form | Administration dosage | Range of application |
|-----------------------|--|---|---|
| Tick . | | | |
| ice | Application is made on 28 m³ empty area. | It is sprayed for 5 - 6 seconds until the surface is wet. | In animal shelters When it's necessary |
| Filea | | | |

PACKAGING: 150 ml SHELF LIFE: 24 months







WELLNESS & GROOMING



BİYO **ICHTHYODERM**



SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g COMPOSITION: 11% Ichthyol.

RECOMMENDED APPLICATION: Biyo Ichthyoderm Pomade is prepared as a skin care product and has emollient, soothing, protective and abscess maturing properties. It enhanced the effect of the main treatment by eliminating the oily crust, dry crust and follicle crusts on the skin caused by various skin diseases of cats, dogs, horses, small and large ruminants. It accelerates the treatment of animals by quickly maturing the abscesses formed. Eliminates itching and removes oily substances (comedones) that fill the sebaceous follicles in the skin. It has mild antiseptic effect against dermatitis. It softens the skin and with this property, it is applied as an emollient on udder skin in mastitis.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin when necessary.

PACKAGING: 10 g, 30 g, 80 g, 100 g SHELF LIFE: 2 years









IYODERM









SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g COMPOSITION: 2% iodine, 10% potassium iodide.

RECOMMENDED APPLICATION: Biyo lyoderm Pomade is a skin care product used for cats, dogs, horses, small and large ruminants. Due to its resolutive and revulsive properties, it helps to remove skin swellings caused by trauma. It has irritant effect. It is used as a protective agent on superficial wound infections, skin irritations, insect bites and small bruises.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin when necessary.

PACKAGING: 10 g, 30 g, 80 g, 100 g SHELF LIFE: 2 years











BİYO **KAMFODERM**









SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g COMPOSITION: 11% camphor

RECOMMENDED APPLICATION: Biyo Kamfoderm is a skin care product used for cats, dogs, horses, small and large ruminants. The camphor in its content is a terpenoid and is a herbal substance obtained from the Cinnamomum camphora plant. Due to its antiseptic, analgesic and resolutive properties, it helps to remove painful muscle swelling and stiffness, itchy skin irritations of horses, small and large ruminants, cats and dogs. It provides natural regeneration and granulation of the skin by covering dry and cracked skin.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin when necessary. Cats and dogs should be prevented from licking the areas where Biyo Kamfoderm has been applied.

PACKAGING: 10 g, 30 g, 80 g, 100 g SHELF LIFE: 2 years









BIYO SALIDERM



Orta Bo 80 g

рy









SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g

COMPOSITION: 15% salicylic acid

RECOMMENDED APPLICATION: Biyo Saliderm Pomade is a cream prepared as a skin care product. It has keratolytic character. It softens keratin, a protein in the structure of the upper layer of the skin, and softens the dried skin cells, hardened and thickened skin layers and allows them to fall off. It helps the main treatment in eliminating formations such as scabs and warts that occur as a result of some skin disorders. It is used to remove dried skin and thickened and hardened skin layers of cattle, sheep, goats, horses, dogs and cats.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin when necessary.

PACKAGING: 10 g, 30 g, 80 g, 100 g

SHELF LIFE: 2 years







BIYO RIVODERM



Büyük Boy 100 g Orta Boy 80 g Küçük Boy 30 g







SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g COMPOSITION: 0.2% Rivanol (Acrinol)

RECOMMENDED APPLICATION: Biyo Rivoderm Pomade is a skin care product. When applied to the sensitive and damaged skin of cats, dogs, horses, small and large ruminants; it acts as a bandage and protects the sensitive and damaged skin from external factors (dirt, moisture, UV, irritating agents, impact, poisons leaking from plants such as ivy, oak tree, sumac). By covering dry and cracked skin, it supports the natural regeneration of the skin, makes it soft and gains elasticity, prevents dehydration of the irritated skin. It accelerates healing by covering mild skin irritations, skin rashes, skin cracks.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin for 3-4 days.

PACKAGING: 10 g, 30 g, 80 g, 100 g **SHELF LIFE:** 2 years







BIYO ZINCODERM



Büyük Boy 100 g

Orta Boy 80 g









SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g

COMPOSITION: 15% zinc oxide

RECOMMENDED APPLICATION: Biyo Zincoderm is a skin care product used for cats, dogs, horses, small and large ruminants. When applied to the sensitive and damaged skin of cats, dogs, horses, small and large ruminants; it acts as a bandage and protects the sensitive and damaged skin from external factors (dirt, moisture, UV, irritating agents, impact, poisons leaking from plants such as ivy, oak tree, sumac). By covering dry and cracked skin, it supports the natural regeneration of the skin, makes it soft and gains elasticity, prevents dehydration of the irritated skin. It accelerates healing by covering mild skin irritations, skin rashes, skin cracks. This product enhances the effect of the main treatment.

USAGE AND DOSAGE: Before using, hairs on the skin should be washed with soap and water and dried. It can be applied 2-3 times a day by rubbing the skin when necessary.

PACKAGING: 10 g, 30 g, 80 g, 100 g **SHELF LIFE:** 2 years







BİYO DERMACURE TRİCOBOR



Büyük Boy 100 g Orta Boy 80 g Küçük Boy 30 q







SKIN CARE POMADE 10 g - 30 g - 80 g - 100 g **RECOMMENDED APPLICATION:** Biyo Dermacure Tricobor is a skin care product used for cats, dogs, horses, small and large ruminants. It has keratolytic and lipolytic character. It softens the keratin in the upper layer of the skin and softens the dried skin cells, hardened and thickened skin layers and allows them to fall off. In addition, with its lipolytic feature: It provides the skin to be degreased and moisturized, provides a good appearance of the skin tissue, and cleans the hair base and follicles It helps the main treatment in eliminating scabs, warts and similar formations that occur as a result of some skin disorders. It supports the natural regeneration of the skin, allowing it to soften and

USAGE AND DOSAGE: Biyo Dermacure Tricobor Pomade, is used for the skincare of body, face, ears, tail, paws and nails of cats, dogs and horses. Before using, hairs on the skin should be cut and washed with soap and water and dried. It is squeezed up to 5 mm into 5 cm² sections on the skin to be applied every day, and it is spread and impregnated. It can be applied 2-3 times a day when necessary

PACKAGING: 10 g, 30 g, 80 g, 100 g **SHELF LIFE:** 3 years

gain elasticity.




COMPOSITION: 2% Chlorhexidine Hydrochloride, 5% Glycolic Acid, 5% Boric Acid.









VERAJEL





....

SKIN CARE PRODUCT 80 g **COMPOSITION:** It contains glycerin, dexpanthenol, aloe vera, potassium sorbate, phenoxyethanol, EDTA.4Na, hydroxy ethyl cellulose, polysorbate 80, purified water.

RECOMMENDED APPLICATION: Verajel is used to support the treatment of wounds caused by various reasons in small and large ruminants, horses, cats, dogs, rabbits, and to relax irritation after itching caused by external parasites such as lice, fleas and ticks. Due to its gel structure and the agents it contains, it covers the wound surface, provides long-term moistening and helps the healing process



USAGE AND DOSAGE: Verajel is applied to the wound surface after mechanical cleaning of the wound surface.

PACKAGING: 80 g SHELF LIFE: 2 years










SHAMPOOS





BIYO DERMACURE TRICOBOR





SHAMPOO 50 ml -100 ml - 250 ml - 450 ml - 1 lt - 5 lt

COMPOSITION: It contains; Chlorhexidine HCI, Glycolic Acid and Boric acid.

RECOMMENDED APPLICATION: It is a skin care product prepared for cats, dogs and horses. It has keratolytic and lipolytic character. It softens the keratin in the upper layer of the skin and softens the dried skin cells, hardened and thickened skin layers and allows them to fall off. In addition, with its lipolytic feature; It provides the skin to be degreased and moisturized, provides a good appearance of the skin tissue, and cleans the hair base and follicles. It helps the main treatment in eliminating scabs, warts and similar formations that occur as a result of some skin disorders. It supports the natural regeneration of the skin, allowing it to soften and gain elasticity.

USAGE AND DOSAGE: The hair of cats and dogs is wetted with water and friction is done. Approximately 2-3 ml of shampoo is spread by foaming. It is applied in the same way by using enough shampoo in horses. After waiting for 5-10 minutes, rinse with plenty of water and dry.

PACKAGING: 50 ml, 100 ml, 250 ml, 450 ml, 1 lt, 5 lt **SHELF LIFE:** 3 years





BIYO DERMACURE **HERBIO**





DRY FOAM SHAMPOO FOR DOGS AND CATS 250 ml

COMPOSITION: It is a colorless, coconut scented shampoo. It contains Palmarosa oil, Ammonium lauryl sulfate, Cocamidopropyl betaine, Cocamido DEA, Polysorbate 20, Phenonip, Aloe vera, Citric acid, Coconat fragrans.

RECOMMENDED APPLICATION: It is a foam shampoo for cats and dogs to be used without water. It contains natural Palmarosa oil obtained from the Cymbopogon martinii plant. Palmarosa oil, which is used in alternative treatment in the East, especially in India, ensures that the skin and hair of cats and dogs are healthy and shiny with its moisturizing properties. It protects the natural oil balance of the skin and prevents skin from become infected.

The foam prepared with a special method cleanses the skin deeply with the cleansing ingredients in its composition and makes the hair look healthier by giving shine to the hair. It gives a pleasant and lasting scent because of the essence in its content. It doesn't irritate the eyes of the animal and does not cause tears.

USAGE AND DOSAGE: It is sprayed into the palm of the hand in the form of foam with its special pump. Starting from the nape of the neck, it is frictioned with the hand to ensure that it penetrates the skin and hair well. The foams are left on the animal for 5-10 minutes. The animal is wiped dry with a towel and combed.

PACKAGING: 250 ml SHELF LIFE: 2 years



BIYO DERMACURE **CURLY**





FOR CURLY CATS & DOGS

250 ml

COMPOSITION: It is a violet-purple shampoo with a lavender scent which contains Polyquarternium-39, PEG-200 Hydrogenated Glyceryl Palmate, PEG-7 Glyceryl Cocoate, Cocamidopropyl Betaine, Sodium Cocoamphoacetate, Disodium Laureth Sulfosuccinate, Decyl glucoside (and) Xylose (and) Decyl alcohol, Cocamido MIPA&laureth-4, Silicone Quarternium-17 (and) Trideceth-7 (and) Trideceth-5 (and) Methylchloroisothiazolinone (and) Methylisothiazolinone, Phenonip, D&C Violet No:2, Fragrance, Citric Acid.

RECOMMENDED APPLICATION: It softens the hardened and curly hairs of dogs and cleans the hairs deeply. It allows the tangled hairs to be opened and brushed easily. It gives shine to the hair and makes it looks healthier. The essence does not only give a pleasant and long-lasting fragrance, it also makes the washing process easier. It is used for cleaning hard, thick and curly haired dogs over 6 weeks old.

USAGE AND DOSAGE: The hair of dogs is wetted with water and friction is done. Approximately 10 ml of shampoo is spread by foaming. After waiting for 3-5 minutes, rinse with plenty of water and dry. If necessary, the application is repeated.

PACKAGING: 250 ml SHELF LIFE: 2 years



BIYO DERMACURE **FRESH**

DEODORANTLI KÖPEK BAKIM SAMPUANI










DEODORANT SHAMPOO

COMPOSITION: It is a lavender-scented shampoo of blue-violet color which contains Peg-200 Hydrogenated Clyceryl Palmate & PEG-7 Glyceryl Cocoate, Cocamidopropyl Betaine, Sodium Cocoamphoacetate, Disodium Laureth Sulfosuccinate, Decyl Glucoside (and) Xylose (and) Decyl Alcohol (and) Water, Cocamide MIPA & Laureth-4, Silicone Quaternium-17 (and) Trideceth-7 (and) Trideceth-5 (and) Methylchloroisothiazolinone (and) Methylisothiazolinone, Sodium Bicarbonate, Phenonip, FD&C Blue No:1, Ext.D&C Violet No:2, Fragrance, Citric Acid.

RECOMMENDED APPLICATION: It deeply cleans the coat due to the cleansing substances. It gives a fresh and clean smell and softens coat and allows them to be brushed comfortably. It gives shine to the coat and makes them look healthier. The product is for dogs over 6 weeks old.

USAGE AND DOSAGE: The hair of dog is wetted with water and friction is done. Approximately 10 ml of shampoo is spread by foaming. After waiting for 3-5 minutes, rinse with plenty of water and dry.

PACKAGING: 50 ml, 100 ml, 250 ml, 450 ml, 1 lt, 5 lt SHELF LIFE: 2 years



50 ml - 100 ml - 250 ml - 450 ml - 1 lt - 5 lt

BİYO DERMACURE **WHİTE**





SKIN CARE SHAMPOO FOR WHITE DOGS 50 ml - 100 ml - 250 ml - 450 ml - 1 lt - 5 lt

COMPOSITION: It is a white-colored, coconut-scented shampoo which contains Peg-200, Hydrogenated Glyceryl Palmate & Peg-7 Glyceryl Cocoate, Cocamidopropyl Betaine, Sodium Cocoamphoacetate, Disodium Laureth Sulfosuccinate, Decyl Glucoside (and) Dodecyl Alcohol (and) Water, Cocamide MIPA & Laureth-4, Silicone Quaternium-17 (and) Trideceth-7 (and) Trideceth-5 (and) Methylchloroisothiazolinone (and) methylisothiazolinone, Sodium Bicarbonate, Anionic, 44'-bis-(Sulfostyryl)-Biphenyl Disodium, Alpha Tridecyl Omega Hydroxypoly (oxy-1,2-Ethanediyl) and Sodium Dodecyl Sulphate, Phenonip, Coconut Fragrance, Citric Acid.

RECOMMENDED APPLICATION: The product α is cleaning and optical whitening substances which cleans and whitens the hair of white dogs deeply. It makes the hair look healthier and prevents the hairs from turning yellow throughout time. It gives a pleasant and long-lasting fragrance due to the essence in its content. It is used for cleaning white dogs older than 6 weeks.

USAGE AND DOSAGE: The hair of dog is wetted with water and friction is applied. Approximately 10 ml of shamped is spread by foarning. After waiting for 3-5 minutes, rinse with plenty of water and dry. If necessary, the application is repeated. Please avoid contact with eyes.

PACKAGING: 50 ml, 100 ml, 250 ml, 450 ml, 1 lt, 5 lt

SHELF LIFE: 2 years



BIYO DERMACURE **BLACK**





SKIN CARE SHAMPOO FOR BLACK DOGS 250 ml

COMPOSITION: It is a black-colored, coconut-scented shampoo which contains PEG-200 Hydrogenated Glyceryl Palmate, PEG-7 Glyceryl Cocoate, Cocamidopropyl Betaine, Sodium Cocoamphoacetate, Disodium Lureth Sulfosuccinate, Dodecyl Glucoside (and) Xylose (and) Decyl Alcohol, Cocamide MIPA&laureth-4, Silicone Quarternium-17 (and) Trideceth-7 (and) Tridecedth-5 (And) Methylchloroisothiazolinone, Glycol Distearat, Sodium Bicarbonate, Phenonip, Fragrance, Citric Acid.

RECOMMENDED APPLICATION: Due to its special color protection feature, it preserves the original color tones of dogs with dark hair such as black, mahogany and chocolate. It makes their hair look healthier by giving shine. It gives a pleasant and long-lasting fragrance due to the essence in its content. It is used for cleaning black dogs older than 6 weeks.

USAGE AND DOSAGE: The hair of dog is wetted with water and friction is done. Approximately 10 ml of shampoo is spread by foaming. After waiting for 3-5 minutes, rinse with plenty of water and dry.

PACKAGING: 250 ml SHELF LIFE: 2 years



BIYO DERMACURE COMFORT





SKIN CARE SHAMPOO FOR CATS & DOGS 250 ml


COMPOSITION: It is a shampoo with a green color and a lavender fragrance which contains in 1 g: 0,01 g Pramoxine HCl, Sodium lauryl glucose carboxylate (and) lauryl glucoside, Coco glucoside (and) Clyceryl oleate, Sodium Lauryl Ether Sulfate.2moles EO, Poliquarternium-7, Cocamide DEA, Polyoxyethylene 20 Sorbitan Monooleate, Fragrance, Colloidal oatmeal, Nipagin M sodium, Nipasol sodium, D & C Green No: 5.

RECOMMENDED APPLICATION: It is used in cats and dogs to reduce skin irritation caused by Pruritus medical itching, which is a symptom of other diseases, and to relieve the animal. In cats and dogs, allergic diseases, ectoparasites, bacterial infections or idiopathic conditions cause itching. In this case, the animal's self-scratching increases skin irritation. When treating the cause of itching, Biyo Dermacure COMFORT Shampoo is used to assist the treatment and prevent itching.

It gives relaxation to your pet, while the colloidal oatmeal it contains gives shine and vitality.

USAGE AND DOSAGE: The hair of cats and dogs is wetted with water and friction is done. Approximately 5-10 ml of shampoo is spread by foaming. After waiting for 3-5 minutes, rinse with plenty of water and dry.

PACKAGING: 250 ml SHELF LIFE: 2 years



BIYO DERMACURE **PUPPY**






50 ml - 100 ml - 250 ml - 450 ml - 1 lt - 5 lt

COMPOSITION: It is a clear orange-colored, lavender-scented shampoo which contains Texapon ASV 50 (Sodium Laureth Sulfate, Sodium Laureth S-Sulfate, Magnesium Laureth Sulfate, Magnesium Laureth S-Sulfate, Sodium Oleth Sulfate, Magnesium Oleth Sulfate), Plantapon LGS Sorb (Sodium Lauryl Glucose Carboxylate, Lauryl Glucoside), Dehyton MC (Sodium Cocoamp hoacetate), Lamesoft PO65 (CocoGlucoside, Glyceryl Oleate), Emulgin SML 20, Arlypon TT (PEG/PPG 120/10 Trimethylolpropane Trioleate, Laureth-2), Dehyquart 701/N (Polyquarternium-7), Phenonip, Glycerin, Wheat Proteine, Aloe vera, D&C Red No. 33, D&C Yellow No. 10, EDTA Disodium, Citric Acid, Fragrance.

RECOMMENDED APPLICATION: It cleans the skin with the cleansing agents and wheat protein in its content, makes the hair look healthier by giving shine. It doesn't irritate the puppy's eyes, does not cause tears. Due to the essence in its content, it gives a pleasant and lasting fragrance and also calming properties of the essence, the washing process becomes easier. It is used for cleaning puppies over 6 weeks old.

USAGE AND DOSAGE: The hair of dog is wetted with water and friction is done. Approximately 10 ml of shampoo is spread by foaming. Please avoid contact with eyes. After waiting for 3-5 minutes, rinse with plenty of water and dry. If necessary, the application is repeated.

PACKAGING: 50 ml, 100 ml, 250 ml, 450 ml, 1 lt, 5 lt **SHELF LIFE:** 2 years



 $\overline{}$

PET-DUST DRY POWDER



1,40





DRY SHAMPOO FOR CATS AND DOGS 100 gr - 500 gr **COMPOSITION:** It is a dry shampoo for skin care which contains Zinc oxide, Boric Acid, Kaolin, Magnesium Stereate and Lavender Aroma.

RECOMMENDED APPLICATION: It is a dry shampoo for cats and dogs. Cats and dogs that need to be washed frequently may experience deterioration in the oil balance of their skin. Such situations deter cats and dogs to be washed with water. Pet dust dry powder ensures the skin is cleaned and removes bad odors and leave a pleasant smell thereafter. This product has the benefit of easy application compared to wet washing. The product is especially advantageous when traveling.

USAGE AND DOSAGE: Massage with the shampoo into dry fur avoiding eyes and mouth. After the excess is cleaned with the help of a dry cloth, the hairs are combed. The animal is dried with a towel. There is no harm in using it frequently.

PACKAGING: 100 g, 500 g **SHELF LIFE:** 3 years



BIYO DERMACURE HERBIO DRY FOAM





COMPOSITION: It is a colorless, coconut scented shampoo. It contains Palmarosa oil, Ammonium lauryl sulfate, Cocamidopropyl betaine, Cocamido DEA, Polysorbate 20, Phenonip, Aloe vera, Citric acid, Coconat fragrans.

RECOMMENDED APPLICATION: It is a foam shampoo for cats and dogs to be used without water. It contains natural Palmarosa oil obtained from the Cymbopogon martinii plant. Palmarosa oil, which is used in alternative treatment in the East, especially in India, ensures that the skin and hair of cats and dogs are healthy and shiny with its moisturizing properties. It protects the natural oil balance of the skin and prevents skin from b ecome infected. The foam prepared with a special method cleanses the skin deeply with the cleansing ingredients in its composition and makes the hair look healthier by giving shine to the hair. It gives a pleasant and lasting scent because of the essence in its content. It doesn't irritate the eyes of the animal and does not cause tears.


USAGE AND DOSAGE: It is sprayed into the palm of the hand in the form of foam with its special pump. Starting from the nape of the neck, it is frictioned with the hand to ensure that it penetrates the skin and hair well. The foams are left on the animal for 5-10 minutes. The animal is wiped dry with a towel and combed.

PACKAGING: 250 ml. SHELF LIFE: 2 years

120



BIYO DERMACURE DRY FOAM





COMPOSITION: It is a colorless, vanilla-scented shampoo which contains Ammonium lauryl sulfate, Cocamidopropyl betaine, Cocamide DEA, Polysorbate 20, Phenonip, Aloe vera, Citric acid, Vanilla fragrance

RECOMMENDED APPLICATION: It is a foam shampoo for cats and dogs, intended for use without water. The foam prepared by a special method cleanses the skin deeply with the cleansing substances in its composition and gives shine to the coat, making them look healthier. It gives a pleasant and lasting smell. It does not irritate eyes, does not cause tears.

USAGE AND DOSAGE: Massage with the shampoo into dry fur avoiding eyes and mouth. The foams are left on the animal for 5-10 minutes. The animal is dried with a towel.

PACKAGING: 250 ml SHELF LIFE: 2 years







BIYO DERMACURE CAT-LINE









SOLUTION 1 ml

COMPOSITION: Tea Tree Oil, Cinnamon Oil, Neem Oil and Lavender Oil

RECOMMENDED APPLICATION: It is a skin care product for cats. With the moisturizing effect of the essential oils it contains, the solution helps to maintain the Ph/moisture balance of the skin without causing the skin's natural oil to be lost, and provides a soft, bright and natural vitality without irritating the skin. It prevents the skin from becoming infected. It is suitable for use on skin that has an allergic reaction to flea, tick and lice bites. It relieves skin irritations caused by fleas, ticks and lice. It enhanced the effect of the main treatment by eliminating crusts, warts and similar formations that occur as a result of some skin disorders. Without affecting the cat's normal flora, supplies a good smell by removing the bad odor.

USAGE AND DOSAGE: It is applied topically directly on the skin. The hairs between the shoulders are opened so that the skin is visible. It is applied in the line between the shoulders. It is appropriate to use 1 microtube for every 15 days. It should not be used in cats under 8 weeks of age.

PACKAGING: 1 ml. SHELF LIFE: 2 years





BIYO DERMACURE DOG-LINE



SOLUTION

1 ml - 2 ml - 4 ml

COMPOSITION: Tea Tree Oil, Cinnamon Oil, Neem Oil and Lavender Oil.

RECOMMENDED APPLICATION: It is a skin care product for dogs. With the moisturizing effect of the essential oils it contains, the solution helps to maintain the Ph/moisture balance of the skin without causing the skin's natural oil to be lost, and provides a soft, bright and natural vitality without irritating the skin. It prevents the skin from becoming infected. It is suitable for use on skin that has an allergic reaction to flea, tick and lice bites. It relieves skin irritations caused by fleas, ticks and lice. It enhanced the effect of main treatment in eliminating crusts, warts and similar formations that occur as a result of some skin disorders. Without affecting the normal flora of the dog, it supplies a good smell by removing the bad odor.

USAGE AND DOSAGE:It is applied topically directly on the skin. The hairs between the shoulders of an upright dog are opened so that the skin can be seen. It is applied in the line between the shoulders. It is appropriate to use 1 microtube for every 15 days. It should not be used in dogs under 8 weeks of age.

PACKAGING: 1 ml, 2 ml, 4 ml SHELF LIFE: 2 years



BIYO DERMACURE HERBIO CATSPRAY



1,48





PUMP LIQUID SPRAY 50 ml - 100 ml - 150 ml - 200 ml

COMPOSITION: 78.79 mg/ml Palmarosa oil.

RECOMMENDED APPLICATION: Herbio Catspray contains natural Palmarosa oil derived from the Cymbopogon martinii plant. Palmarosa oil, which is used in alternative medicine in the Far East, especially in India, provides the skin and hair of cats to be healthy and shiny with its moisturizing feature. It maintains the natural oil balance of the skin. It prevents the skin from becoming infected.

It also helps to treat cat wounds and bruises, and relieves skin irritations caused by fleas, ticks and lice.

USAGE AND DOSAGE: It is sprayed from a distance of 15 cm until all the hairs become wet. The hairs are brushed in the direction of their growth. It can be repeated after 15 days if necessary.

PACKAGING: 50 ml, 100 ml, 150 ml, 200 ml SHELF LIFE: 2 years









120



PUMP LIQUID SPRAY 50 ml - 100 ml - 150 ml - 200 ml


COMPOSITION: 78.79 mg/ml Palmarosa oil.


RECOMMENDED APPLICATION: Herbio Dogspray contains natural Palmarosa oil derived from the Cymbopogon martinii plant. Palmarosa oil, which is used in alternative medicine in the Far East, especially in India, provides the skin and hair of dogs to be healthy and shiny with its moisturizing feature. It maintains the natural oil balance of the skin. It prevents the skin from becoming infected. It also helps to treat dog wounds and bruises, and relieves skin irritations caused by fleas, ticks and lice.

USAGE AND DOSAGE: It is sprayed from a distance of 15 cm until all the hairs become wet. The hairs are brushed in the direction of their growth. It can be repeated after 15 days if necessary.

PACKAGING: 50 ml, 100 ml, 150 ml, 200 ml







VERABLU



المعالم المعال

AEROSOL SPRAY

SKIN CARE PRODUCT

200 ml

COMPOSITION: It contains 40 mg Brilliant Blue in 100 g as an active ingredient and Glycerin, Aloe Vera, Vitamin E, Vitamin A, Dexpanthenol, Isopropyl Alcohol, LPG as excipient.

RECOMMENDED APPLICATION: It is an epithelial protective and physical concealer to protect the area from external effects in lesions on the skin and nails and to help natural healing in cattle, sheep, goats, horses, cats and dogs.

USAGE AND DOSAGE: The lesioned area is cleaned. It is applied to the cleaned area by spraying 1-2 times a day for 1-2 seconds.

PACKAGING: 200 ml SHELF LIFE: 2 years









100 g - 250 g - 500 g - 1000 g - 17 kg

COMPOSITION: Each 100 g contains 15 g phenol, 50 g medical soft soap, 20 g pine tar, 15 g isopropyl alcohol.

RECOMMENDED APPLICATION: It is used for nail care of large and small ruminants and horses.

USAGE AND DOSAGE: The Kreolin Solution should be diluted 5% (approximately 1000 g per 1 tin of water) and the feet of the animals are kept in the resulting emulsion for 1-2 minutes 2-3 times a day. It can be applied to the cuticle of the animals with a brush.

PACKAGING: 100 g, 250 g, 500 g, 1000 g, 17 kg **SHELF LIFE:** 2 years







BORDUST






COMPOSITION: BORDUST contains Tea Tree Oil, Cinnamon Oil, Eucalyptus Oil, Lavender Oil, Zinc Oxide and Boric Acid.


RECOMMENDED APPLICATION: BORDUST is a skin care product prepared for use on ovine and bovine animals, cats, dogs, horses and poultry. With the moisturizing effect of the essential oils it contains, it maintains the pH/moisture balance of the skin without causing the skin to lose its natural oil. It protects the skin against the harmful effects of external factors such as flea, lice and tick bites, sun rays, bad odors and toxins and relieves skin irritations. BORDUST, which helps the main treatment in eliminating scabs, warts and similar formations that may occur on the skin, helps to protect the skin health of animals.

USAGE AND DOSAGE: BORDUST; After sprinkling on the skin of the animals, it should be ensured that the powder is distributed between the hairs on the skin by friction and massage with the help of a brush or by hand.

PACKAGING: 100 g, 250 g, 500 g, 1000 g **SHELF LIFE:** 2 years







BIYO DERMACURE EAR CLEANER





COMPOSITION: Biyo Dermacure Ear Cleaner Solution contains; propylene glycol, salicylic acid, boric acid, acetic acid, glycerine, isopropyl alcohol, and preservative as benzalkonium chloride.

RECOMMENDED APPLICATION: Biyo Dermacure Ear Cleaner Solution, cleans the earwax caused by excessive ear discharge of cats and dogs, delays the formation of dirt, helps to prevent ear infections by preventing necrotic tissues and itching, keeps the ear canal dry and prevents odor.

USAGE AND DOSAGE: Biyo Dermacure Ear Cleaner Solution, as 5-8 drops are poured into the outer ear, after waiting for a while, the base of the ear canal is rubbed to remove the earwax and necrotic tissues. The ear is carefully cleaned with a cotton swab. For very dirty ears, application has to be repeated 2-3 times a day. Routine usage as 2-3 times a week.

PACKAGING: 50 ml, 100 ml SHELF LIFE: 3 years





BIYO DERMACURE EYE CLEANER



EYE CARE SOLUTION - FOR CATS AND DOGS 50 ml - 100 ml **COMPOSITION:** Biyo Dermacure Eye Cleaner Solution contains; boric acid, borax, sodium chloride and preservative as benzalkonium chloride.

RECOMMENDED APPLICATION: Biyo Dermacure Eye Cleaner Solution, cleans the accumulations of eye crust around eyes caused by excessive eye discharge of cats and dogs, delays the formation of dirt and also removes the stains caused by the eye discharge of light-colored dogs and cats.


USAGE AND DOSAGE: Biyo Dermacure Eye Cleaner Solution, applied to each eye as approximately 2.5 ml in every day with the help of cotton, cotton swab or gauze moistened, prevents dirt and accumulation of dirt around the eyes and also removes the stains.

PACKAGING: 50 ml, 100 ml SHELF LIFE: 3 years

120











SPRAY SOLUTION

ORAL AND DENTAL CARE PRODUCT

30 ml - 50 ml - 100 ml -250 ml - 500 ml


COMPOSITION: 0.12% Chlorhexidine Gluconate, 0.20% Aloe Vera and 2% Glycerin

RECOMMENDED APPLICATION: With the Chlorhexidine Gluconate it contains, it is effective in cleaning the oral mucosa in cats and dogs, refreshing the breath and eliminating bad odors in the mouth. It also protects the oral mucosa from harmful factors by forming a protective layer in the mouth. It helps to prevent the formation of plaque and tartar, which causes dental diseases, tooth decay and tooth loss in cats and dogs.

USAGE AND DOSAGE: It can be applied to cats and dogs after each meal or 1-2 times a day. The upper lip is gently lifted to expose the animal's teeth and gingiva. It can be applied as a spray along the gingiva line. During application, care should be taken that the product does not get into the oral cavity of the animal or that the animal does not swallow the product. The product can be applied to animals that are moving or likely to swallow, by wiping the teeth and gingiva with a cotton.

PACKAGING: 30 ml, 50 ml, 100 ml, 250 ml, 500 ml **SHELF LIFE:** 2 years







100 ml

COMPOSITION: It contains acetyl cedrene, ethoxy-methylcyclo-decylether, L-alpha-pinene, isoeugenol, ß-pinene, isoeugenol, ß-pinene, isopropyl alcohol.

RECOMMENDED APPLICATION: Biyo Trainer Spay is a product which designed to help kittens and adult cats and dogs develop the habit of going to the toilet in designated areas (inside or outside the house). The animal will learn to go to the toilet in the designated place after a while due to the smell of the product and will gain a permanent habit.

USAGE AND DOSAGE: Shake the product bottle before use.

INSIDE THE HOUSE: A training pad is laid on the place designated as the toilet area. From a distance of 20-30 cm, spray 4-5 dega of Biyo Trainer Spray over the entire surface and let the animal smell it.


OUTSIDE THE HOUSE: If the toilet area is designated outside the house, this training starts inside the house. A training pad is laid on the place designated as the toilet area. From a distance of 20-30 cm, spray 4-5 dega of Biyo Trainer Spray over the entire surface and let the animal smell it. The application should be continued until the animal gains the toilet habit.

PACKAGING: 100 ml SHELF LIFE: 2 years





SUPPLEMENTS FOR PETS









MULTIVITAMIN TABLET FOR DOGS 60 Tablets

RECOMMENDED APPLICATION: It is used for the vitamin needs of puppies, adult and pregnant dogs. The tablets should be added their

For puppies and small breed dogs : 100 g meal /½ tablet / day. For adult and pregnant dogs :150 g meal /1 tablet / day. : 250 g meal /1+ ½ tablet / day. Large breed dogs

| Active material | Level in premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Vitaminler | | |
| Vitamin A/ Vitamin A Acetate | 8.000.000 | IU/kg |
| Vitamin D ₃ | 300.000 | IU/kg |
| Vitamin E/α-tocopherol acetate | 40.000 | mg/kg |
| Vitamin C/Ascorbic acid | 11.000 | mg/kg |
| Vitamin B ₁ /Thiamine HCl | 2.000 | mg/kg |
| Vitamin B ₂ /Riboflavin-5-phosphate | 2.800 | mg/kg |
| Vitamin B ₃ /Niacinamide | 30.000 | mg/kg |
| Vitamin B ₅ /Calcium-d-pantothenate | 4.000 | mg/kg |
| Vitamin B ₆ /Pyridoxine HCL | 2.000 | mg/kg |
| Vitamin B ₇ /d-Biotin 2% | 40 | mg/kg |
| Vitamin B ₉ /Folic acid | 200 | mg/kg |
| Vitamin B ₁₂ /Siocobalamin | 0,02 | mg/kg |
| Vitamin K ₃ /Menadione sodium bisulfite | 3.000 | mg/kg |
| Sensory additives | | |
| Aromatic mixtures | 30.000 | mg/kg |
| Carriers | | |
| Lactose/Lactose Monohydrate | 806.400 | mg/kg |
| Magnesium Stearate | 10.000 | mg/kg |

PACKAGING: 60 tablets SHELF LIFE: 24 months



BIYO POWERCURE CAT-VIT ONE-A-DAY



MULTIVITAMIN TABLET FOR CATS
60 Tablets


For Adult and Pregnant Cats: 1 tablet/100 g food per day Vitamins A

It is use for the vitamin needs of kittens, adult and pregnant cats. It reduces the breakage and shedding of the hairs and makes them shiny. Increases milk yield of offspring females. It helps kittens to grow quickly.

PACKAGING: 60 tablets SHELF LIFE: 24 months

120

....





BIYO POWERCURE CALCIVIT-D ONE-A-DAY



CALCIUM-PHOSPHORUS-D₃ FOR CATS AND DOGS60 Tablets

RECOMMENDED APPLICATION:

For kittens and adult cats 100g food /1 tablet per day
For pregnant, lactating and elderly cats 150g food /2 tablets per day
For puppies and small breed dogs/150g food 2 tablets per day
For pregnant, lactating and elderly dogs 200g food /3 tablets per day
For large breed dogs 300-500g food /3-5 tablets per day

It meets the calcium, phosphorus and vitamin D_3 needs of puppies, adults and pregnant cats and dogs. It contributes to the strength of bone structures.

| Active material | Level in premix | Units |
|-----------------------------|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin D ₃ | 125.000 | IU/kg |
| Sensory additives | | |
| Aromatic mixtures | 100.000 | mg/kg |
| Carriers | | |
| Lactose/Lactose Monohydrate | 341.040 | mg/kg |
| Magnesium Stearate | 10.000 | mg/kg |
| D.C.P/Dicalcium Phosphate | 555.000 | mg/kg |
| Phosphorus 18 % | | |
| Calcium 25 % | | |



PACKAGING: 60 tablets
SHELF LIFE: 24 months





BREWERS YEAST ONE-A-DAY











PACKAGING: 60 tablets

SHELF LIFE: 24 months

PREMIX GARLIC YEAST TABLET
FOR CAT AND DOGS
60 Tablets

RECOMMENDED APPLICATION: Biyo Powercure Brewers Yeast tablet enriched with garlic powder is a source of protein, trace minerals and B-complex vitamins. It stimulates the appetite of cats and dogs, provides vitamin supplements, reduces hair loss and makes hair shiny, protects the gastrointestinal microflora and increases milk yield of lactating females.

USAGE DESCRIPTION:

| Kittens | 50 g meal/food | 1 Tablet / Day |
|------------------------------|-----------------------|-------------------|
| Adult Cats | 100 g meal/food | 2 Tablets / Day |
| Puppies and Small Breed Dogs | 100 g-200 g meal/food | 2 Tablets / Day |
| Large Breed Dogs | 300 g-500 g meal/food | 3-5 Tablets / Day |

| Active material | Level in | Units |
|---|----------|-------|
| Nutritional Additives | Premix | |
| Vitamins | | |
| Vitamin E/α-Tocopherol Acetate | 12.000 | mg/kg |
| Vitamin C/Ascorbic Acid | 170 | mg/kg |
| Vitamin B2/Thiamine Monohydrate | 370 | mg/kg |
| Vitamin B3/Niacinamide | 920 | mg/kg |
| Vitamin B5/Calcium-Pantetonate | 735 | mg/kg |
| Vitamin B6/Pridoxine HCL | 35 | mg/kg |
| Vitamin B9/Folic Acid | 15 | mg/kg |
| Vitamin B12/Cyanocobalamin | 7,5 | mg/kg |
| Trace elements | | |
| Iron/Iron Sulfate Monohydrate 30% | 1.670 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 185 | mg/kg |
| Manganese/Manganese Sulfate Monohydrate | 370 | mg/kg |
| Zinc/Zinc Sulfate Heptahydrate | 2750 | mg/kg |
| Sensory additives | | |
| Inactive Brewer's Yeast | 830.000 | mg/kg |
| Garlic Powder | 10.00 | mg/kg |
| Aromatic mixtures | 30.000 | mg/kg |
| Carriers | | |
| Lactose/Lactose monohydrate | 70.000 | mg/kg |
| Magnesium Stearate | 20.000 | mg/kg |



METHIO
ONE-A-DAY




AMINO ACID AND VITAMIN TABLET
FOR CATS AND DOGS
60 Tablets

RECOMMENDED APPLICATION: METHIO Tablet is a premix that meets the protein needs of cats and dogs and is fortified with amino acids, minerals and vitamins.

| 50 g food for kittens | 0.5 tablets per day | |
|----------------------------------|-------------------------|--|
| 100 g food for adult cats | 1 tablet per day | |
| 100-200 g of food for dogs up to | 10 kg/1 tablet per day | |
| 200-300 g of food for dogs up to | 20 kg/2 tablets per day | |
| 300-500 g of food for dogs up to | 30 kg/3 tablets per day | |

| Active material | Level in premix | Units | |
|--|-----------------|-------|--|
| Nutritional supplements | | | |
| Vitamins | | mg/kg | |
| Vitamin E | 26.700 | mg/kg | |
| Vitamin C/L-ascorbic Acid Monophosphate | e 21.330 | mg/kg | |
| Vitamin B ₁ /Tiamine Mononitrate | 6.667 | mg/kg | |
| Vitamin B ₂ /Riboflavin | 6.667 | mg/kg | |
| Vitamin B ₅ /Calcium-d-pantothenate | 10.000 | mg/kg | |
| Vitamin B ₆ /Pyridoxine HCL | 6.667 | mg/kg | |
| Vitamin B ₉ /Folic acid | 4 | mg/kg | |
| Vitamin B ₁₂ /Siocobalamin | 13,4 | mg/kg | |
| Vitamin K ₃ /Menadione Sodium Bisulfite | 667 | | |
| Trace elements | | mg/kg | |
| Zinc/Zinc Oxide | 4.000 | | |
| Aminoacids | | mg/kg | |
| DL-Methionine | 333.400 | | |
| Sensory feed additives | | mg/kg | |
| Aromatic mixtures | 7.350 | | |
| Feed ingredients | | mg/kg | |
| Lactose | 182,6346 | mg/kg | |
| Magnesium Stearate | 27,300 | mg/kg | |
| Sorbitol | 366.600 | | |
| | | | |







BİYOFLEX ONE-A-DAY



JOINT HEALTH SUPPLEMENT FOR CATS AND DOGS 30 Tablets Biyoflex One-A-Day tablet contains Glucosamine Sulfate, Chondroitin Sulfate and Methyl Sulfonyl Methane. BİYOFLEX One-A-Day tablet protect the physiological structure of the joints and cartilage tissue of cats and dogs.

RECOMMENDED APPLICATION:

| Body weight | Initial Dose for 4-6 weeks | After 6 weeks |
|-------------------------|----------------------------|-----------------------------|
| Dog and cat up to 8 kg | Half tablet/day | Half tablet/every other day |
| Dog and cat up to 15 kg | 1 tablet/day | Half tablet/day |
| Dog and cat up to 30 kg | 2 tablets/day | 1 tablet/day |
| More than 30 kg dog | 3 tablets/day | 2 tablets/day |

The product is mixed into 100 g of food or you can give directly.

| Active material | Level in premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Trace minerals | 2.800 | |
| Manganese (Manganese Sulfate Tetrahydrate) | | mg/kg |
| Sensory Additives | 100.000 | |
| Aromatic Blends | | mg/kg |
| Active materials | 333.000 | |
| Glucosamine Sulfate | 167.000 | mg/kg |
| Chondroitin Sulfate | 167.000 | mg/kg |
| Methyl Sulfonyl Methane | | mg/kg |
| Carriers | 174.800 | |
| Lactose/Lactose Monohydrate | 30.000 | mg/kg |
| Magnesium Stearate | | mg/kg |

PACKAGING: 30 tablets SHELF LIFE: 24 months





Analytical Components

% 0,014 % 0

% O

% 14

Crude Protein

Crude Cellulose Crude Oil

Crude Ash

BİYOTAN
ONE-A-DAY



VITAMIN TABLET FOR CATS AND DOGS
60 Tablets

RECOMMENDED APPLICATION: It meets the d-biotin needs of cats and dogs. Reduces hair breakage and hair loss and makes them shiny.


USAGE AND DOSAGE:

Kittens and Adult Cats
Puppies and Small Breed Dogs
Medium Breed Dogs
Large Breed Dogs
For Large Breed Dogs

100 g food/1 tablet per day 150 g /food 2 tablets per day 200 g food/3 tablets per day 300-500 g /food 3-5 tablets per day 300-500 g/ food 3-5 tablets per day

| Active material | Level in premix | Units |
|-------------------------------------|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin B ₇ /d-Biotin 2% | 15.000 | mg/kg |
| Sensory Additives | | |
| Aromatic mixtures | 250.000 | mg/kg |
| Carriers | | |
| Lactose/Lactose Monohydrate | 378.000 | mg/kg |
| Magnesium Stearate | 40.000 | mg/kg |
| Sorbitol | 182.000 | mg/kg |

PACKAGING: 60 tablets SHELF LIFE: 24 months





65

64____

BIYO POWERCURE

CAT-FORMULA











PACKAGING: 200 g

SHELF LIFE: 24 months

COMPLEMENTARY FEED FOR KITTENS 200 g

RECOMMENDED APPLICATION: It is a complementary feed which is prepared to meet the special needs of the sensitive stomachs of kittens which are separated from their mother at an early stage or whose mother is not able to get enough milk or orphaned kittens. It is prepared on the basis of natural breast milk and it contains vitamins, minerals, I-Carnitine and taurine. Especially ideal for kittens up to 3 months old. It can be used during pregnancy and lactation.

USAGE AND DOSAGE: Biyo Powercure Cat Formula powder is mixed with a small amount of water. The remain of the water is added little by little with rapid stirring. Fresh mix should be given at every meal.

| 5g | Every 2 hours - day and night |
|---------|--------------------------------------|
| 10 g | Every 2 hours between 6 and 24 hours |
| 15 g | Every 3 hours between 6 and 24 hours |
| | 4-6 meals during the day |
| 15-20 g | In case of need |
| | 10 g 15 g |

| Vitamins Vitamin A Vitamin D ₃ Vitamin K ₃ /Menadione Sodium Bisulfite Vitamin B ₁ /Tiamine Hydrochloride Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 20.000 2.000 0,2 10 10 80 20 | IU/kg IU/kg mg/kg mg/kg mg/kg mg/kg |
|---|--|-------------------------------------|
| Vitamin A Vitamin D ₃ Vitamin N ₃ /Menadione Sodium Bisulfite Vitamin B ₁ /Tiamine Hydrochloride Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 2.000 0,2 10 10 80 | IU/kg mg/kg mg/kg mg/kg |
| Vitamin D ₃ Vitamin K ₃ /Menadione Sodium Bisulfite Vitamin B ₁ /Tiamine Hydrochloride Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 2.000 0,2 10 10 80 | IU/kg mg/kg mg/kg mg/kg |
| Vitamin K ₃ /Menadione Sodium Bisulfite Vitamin B ₁ /Tiamine Hydrochloride Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 0,2 10 10 80 | mg/kg mg/kg mg/kg |
| Vitamin B ₁ /Tiamine Hydrochloride Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 10 10 80 | mg/kg mg/kg |
| Vitamin B ₂ /Riboflavin Universal Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCl Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 10 | mg/kg |
| Vitamin B ₃ /Niacinamide Vitamin B ₅ /Calcium-D-Pantothenate Vitamin B ₆ /Pridoxine HCI Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 80 | |
| Vitamin B _s /Calcium-D-Pantothenate Vitamin B _s /Pridoxine HCI Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | | mg/kg |
| Vitamin B ₆ /Pridoxine HCI Vitamin B ₇ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 20 | |
| Vitamin B ₂ /D-Biotin 2% Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | | mg/kg |
| Vitamin B ₉ /Folic Acid Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 8 | mg/kg |
| Vitamin B ₁₂ /Cyanocobalamin Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 0,2 | mg/kg |
| Trace minerals Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 2 | mg/kg |
| Iron/Iron Sulfate Heptahydrate 20% Zinc/Zinc Sulphate Monohydrate 35% | 0,06 | mg/kg |
| Zinc/Zinc Sulphate Monohydrate 35% | | |
| • | 90 | mg/kg |
| | 50 | mg/kg |
| Manganese/Manganese Sulfate Monohydra | te 32% 5 | mg/kg |
| Iodine/Potassium Iodide 75% | 1 | mg/kg |
| Sodium Selenite Selenium 45% | 0,1 | mg/kg |
| Aminoacids | | |
| Taurine | 1.000 | mg/kg |
| L-Carnitine L-Tartrate | 400 | mg/kg |
| Carriers | | |
| Milk Protein | 100.000 | mg/kg |
| Soy Protein | 180.000 | mg/kg |
| Whole Milk Powder | 720.000 | mg/kg |

| Analytical Components | |
|------------------------------|--------|
| Crude Protein | % 41 |
| Crude Cellulose | % O |
| Crude Ash | % 5,6 |
| Crude Oil | % 19,7 |

BIYO POWERCURE DOG-FORMULA









COMPLEMENTARY FEED FOR PUPPIES 200 g **RECOMMENDED APPLICATION:** It is a complementary feed which is prepared to meet the special needs of the sensitive stomachs of puppies which are separated from their mother at an early stage or whose mother is not able to get enough milk or orphaned puppies. It is prepared on the basis of natural breast milk and it contains vitamins, minerals, I-Carnitine and taurine. Especially ideal for puppies up to 3 months old. It can be use during pregnancy and lactation.

USAGE AND DOSAGE: Biyo Powercure Dog Formula powder is mixed with a small amount of water. The remain of the water is added little by little with rapid stirring. Fresh mix should be given at every meal.

| | Small breeds | Large breeds | Meal order |
|---------------------|--------------|--------------|--------------------------------------|
| rom birth to 1 week | 5 g | 10 g | Every 2 hours - day and night |
| st and 3rd week | 10 g | 30 g | Every 2 hours between 6 and 24 hours |
| ord and 6th week | 20 g | 125 g | Every 3 hours between 6 and 24 hours |
| After 6th week | 40 g | 225 g | Every 4 hours between 6 and 24 hours |
| n adult dogs | 40 g | 225 g | In case of need |

| addit dogs | 9 | 2239 |
|---|-----------------|-------|
| | | |
| ctive material | Level in premix | Units |
| eed additives | | |
| itamins | | |
| tamin A | 20.000 | IU/kg |
| itamin D ₃ | 2.000 | IU/kg |
| tamin K ₃ /Menadione Sodium Bisulfit | e 0,2 | mg/kg |
| itamin B ₁ /Tiamine Hydrochloride | 10 | mg/kg |
| itamin B ₂ /Riboflavin Universal | 10 | mg/kg |
| itamin B ₃ /Niacinamide | 80 | mg/kg |
| itamin B _s /Calcium-D-Pantothenate | 20 | mg/kg |
| itamin B ₆ /Pridoxine HCI | 8 | mg/kg |
| itamin B ₇ /D-Biotin 2% | 0,2 | mg/kg |
| itamin B ₉ /Folic Acid | 2 | mg/kg |
| itamin B ₁₂ /Cyanocobalamin | 0,06 | mg/kg |
| race elements | | |
| on/Iron Sulfate Heptahydrate 20% | 90 | mg/kg |
| inc/Zinc Sulphate Monohydrate 35% | 50 | mg/kg |
| anganese/Manganese Sulfate Monohyo | drate 32% 5 | mg/kg |
| dine/Potassium Iodide 75% | 1 | mg/kg |
| odium Selenite Selenium 45% | 0,1 | mg/kg |
| minoacids | | |
| aurine | 1.000 | mg/kg |
| Carnitine L-Tartrate | 400 | mg/kg |
| arriers | | |
| ilk Protein | 100.000 | mg/kg |
| by Protein | 180.000 | mg/kg |
| /hole Milk Powder | 720.000 | mg/kg |

Analytical Components

Crude Protein % 41

Crude Cellulose % 0

Crude Ash % 5,6

Crude Oil % 19,7

PACKAGING: 200 g SHELF LIFE: 24 months

BIYO-ANISOL PET 120

VITAMIN - AMINO ACID 30 ml

It is a solution which contains endogenous-exogenous amino acids and essential minerals along with water-soluble vitamins.

Composition: 1 It contains

Vitamins, provitamins and substances with chemically well-defined similar properties

Active material / Additive Name (3a831) Vitamin B₆ (Pyridoxine) 500 mg,

(3a820)Vitamin B₁ (Tiamin) 500mg, (3a 826)Vitamin B₂ (Riboflavin-5 phosphate) 500, (3a84)Vitamin B₅ (Pantothenic acid) 1000 mg, (3a315)Niasinamide 2000 mg, (3a890)Choline chloride 20.000mg, (3a880)Biotin 3 mg

Amino acids Salts and analogs of amino acids: Active material / Additive Name

(3c440)L- Tryptophan 1500 mg, (3c301)DL-Methionine 3000 mg, (3c321) L-Lysine 8000mg, (3c410) L-Threonine 4200 mg,

Flavoring agents

(2b02010)Benzyl alcohol 10.000 mg

Component List

(11.4.1)Sodium chloride 18.000 mg,(11.2.6) Magnesium chloride 170 mg

Analytical Components: Crude Protein 1%, Crude Fat 0%, Crude Cellulose 0%, Crude Ash 1.1%

RECOMMENDED APPLICATION: As a supportive it meets the vitamin and amino acid needs of all target animal species listed below. It is also used as a supplement in accelerating their development, increasing their productivity, increasing feed efficiency and strengthening the immune system during illness.

USAGE AND DOSAGE: It can be used by mixing drinking water of animals or by drinking directly. It should be ensured that animals consume the medicated water within 7-8 hours at least.

Animal species / Dosage

Canary and Budgerigar: 1-2 drops Parrot and Pigeon: 4-5 drops Cat and Dog: 2-3 ml/10 kg b.w

PACKAGING: 30 ml. SHELF LIFE: 24 months











FERMACAT











ANTI - HAIRBALL CAT MALT 80 g

RECOMMENDED APPLICATION: Due to the malt and vegetable oils in its content, it facilitates the natural exit of swallowed hairs and other indigestible components. It doesn't contain artificial dyes and flavoring agent.

USAGE AND DOSAGE: It can be given at the rates specified in the table below, by squeezing directly from the tube into the mouth or by placing it on the paw. It can also be given by adding the meal.

PACKAGING: 80 g SHELF LIFE: 24 months

| JSAGE AIM | Frequency of use | Usage amount |
|------------------------|-----------------------|--------------|
| Excretion of hairballs | Once a day for 5 days | 2,5-5,0 cm |
| Prevent of hairballs | 2-3 times a week | 2,5-5,0 cm |

| Level in premix | Units |
|------------------|--|
| Level III premix | Tilles |
| | |
| | |
| 25.000 | mg/kg |
| 50.000 | mg/kg |
| 50.000 | mg/kg |
| | |
| 500 | mg/kg |
| | |
| 15.000 | mg/kg |
| | |
| | |
| 10.000 | mg/kg |
| | |
| 400.000 | mg/kg |
| 200.000 | mg/kg |
| 30.000 | mg/kg |
| | 50.000 50.000 5000 15.000 10.000 400.000 200.000 |



Crude Ash Moisture % 0,31

% 22,23



FELAXA BİYO-ZINC



ANTI-HAIRBALL AND ZINC-CONTAINING MALT FOR CATS 80 g COMPOSITION: Biotin (3a880) 2000 mg/kg, Zinc sulfate (E6) 1500 mg/kg, Malt extract (1.1.18) 32.8%, Vegetable oils (Palm, 2.20.1) 20%, Yeast (Inactive Brewer's Yeast, 12.1.12) 3%, Liquid Lecithin (1c322) 50.000 mg/kg, Liquid Caramel (E150) 10.000 mg/kg, Glycerin (E422) 49.850 mg/kg

ANALYTICAL COMPONENTS: Crude Protein 6%, Crude Fat 25%, Crude Cellulose 2.5%, Crude Ash 1.3%, Moisture 15%

RECOMMENDED APPLICATION: Biotin helps to slow down hair loss in cats and increase the durability of nails. It accelerates the growth of kittens. Zinc is essential for immune, nervous and reproductive functions and contributes to accelerated wound healing. In addition, malt and vegetable oils facilitate the natural removal of ingested hair and other indigestible components. It doesn't contain artificial dyes or flavors.

USAGE AND DOSAGE: It can be given directly from the tube into the mouth or placed on the paw, or it can be added to the food. For adult cats: 2 cm/day
For kittens: 1 cm/day

PACKAGING: 80 g SHELF LIFE: 2 years







FELAXA JOINT



JOINT - BONE - MUSCLE SUPPORTING MALT 80 g

COMPOSITION: Glucosamine HCL (13.2.8) 300 mg/kg, Msm (methyl sulfonyl methane) (13.9.1) 300 mg/kg-Chondroitin sulfate (13.7.1): 100 mg/kg, Malt extract (1.1.18) 32.8%, Vegetable oils (Palm, 2. 20.1%) 20%, Yeast (Inactive Brewer's Yeast, 12.1.12) 3%, Liquid Lecithin (1c322) 50.000 mg/kg, Liquid Caramel (E150) 10.000 mg/kg, Glycerin (E422) 49.850 mg/kg

ANALYTICAL COMPONENTS: Crude Protein: 8% Crude Fat: 18% Cellulose: 3.0% Crude Ash: 2.1% Moisture: 13%.

RECOMMENDED APPLICATION: In order to support joint health in cases such as aging, obesity, stress, malnutrition in cats, it helps to strengthen cartilage tissue, ligaments and joints, helps to regulate joint fluid in acute or chronic arthritis conditions and helps to increase mobility. In addition, malt and vegetable oils facilitate the natural removal of ingested hair and other indigestible components. It doesn't contain artificial dyes or flavors.

USAGE AND DOSAGE: It can be given directly from the tube into the mouth or placed on the paw, or it can be added to the food. For adult cats: 2 cm/day
For kittens: 1 cm/day

PACKAGING: 80 g SHELF LIFE: 2 years





FELAXA MULTIVITAMIN



MULTIVITAMIN + AMINOACID MALT 80 g

COMPOSITION: Malt extract (1.1.18) 32.8%, Vegetable oils (Palm, 2.20.1) 20%, Yeast (Inactive Brewer's Yeast, 12.1.12) 3%, Vit. B6(3a831) 500 mg/kg, Vit. B1(3a820) 500 mg/kg, Vit. B2(3a826) 500 mg/kg, Vit. B5(3a841) 1.000 mg/kg, Niacinamide(3a315) 2.000 mg/kg, Choline Chloride (3a890) 20.000 mg/kg, Biotin (3a880) 3 mg/kg, L-Triptophane (3c440) 1.500 mg/kg, DL-Methionine (3c301) 3.000 mg/kg, L-Lysine (3c321) 8.000 mg/kg, L-Threonine (3c410) 4.200 mg/kg, Liquid Lecithin (1c322) 50.000 mg/kg, Liquid Caramel (E150) 10.000 mg/kg, Clycerin (E422) 49.850 mg/kg

ANALYTICAL COMPONENTS: Crude Protein 1.5% Crude, Fat 20%, Crude Cellulose 1.5%, Crude Ash 0.3

RECOMMENDED APPLICATION: The cats often lick their bodies to clean themselves. In the process, they swallow hair that is weak, broken or about to break off. The swallowed hairs can form a hairball in the intestines over time, causing blockages in the gastrointestinal tract. B Group vitamins, amino acids, biotin and choline chloride contribute the regular functioning of nerve, muscle and heart functions, slowing hair loss, accelerating growth in kittens, converting food into energy, increasing cell repair, developing the nervous system, skin regeneration and protection, strengthening the immune system, increasing nail durability, supporting the nervous system with healthy skin and hair structure, supporting cell regeneration, especially in older cats, contributing to the survival of cells.

USAGE AND DOSAGE: It can be given directly from the tube into the mouth or placed on the paw, or it can be added to the food. For adult cats: 2-4 cm/day

PACKAGING: 80 g SHELF LIFE: 2 years



FELAXA TAURINE



ANTI-HAIRBALL AND TAURINE-CONTAINING MALT 80 g

COMPOSITION: Taurine (3.a.370) 6.000 mg/kg, Malt extract (1.1.18) 32.8%, Vegetable oils (Palm, 2.20.1) 20%, Yeast (Inactive Brewer's Yeast, 12.1.12) 3%, Liquid Lecithin (1c322) 50.000 mg/kg, Liquid Caramel (E150) 10.000 mg/kg, Clycerin (E422) 49.850 mg/kg

ANALYTICAL COMPONENTS: Crude Protein: 8% Crude Fat: 20% Cellulose: 3.0% Crude Ash: 1.1% Moisture: 13

RECOMMENDED APPLICATION: Taurine, an essential amino acid of vital importance for cats, helps strengthen the heart muscle and supports heart health, protects the integrity of the cells in the retina, contributes to the development of cells in the eye and healthy eye functions, supports rapid development, especially in kittens, plays an important role in the functions of the nervous system and immune system, contributes to a healthy skin and coat structure and supports the digestive system. In addition, malt and vegetable oils facilitate the natural removal of ingested hair and other indigestible components. It doesn't contain artificial dyes or flavors.

USAGE AND DOSAGE: It can be given directly from the tube into the mouth or placed on the paw, or it can be added to the food. For adult cats: 2 cm/day
For kittens: 1 cm/day

PACKAGING: 80 g SHELF LIFE: 2 years



FELAXA LYSINE





COMPOSITION: Lysine (L-Lysine HCI) (3c321) 25.000 mg/kg, Malt extract (1.1.18) 32.8%, Vegetable oils (Palm, 2.20.1) 20%, Yeast (Inactive Brewer's Yeast, 12.1.12) 3%, Liquid Lecithin (1c322) 50.000 mg/kg, Liquid Caramel (E150) 10.000 mg/kg, Glycerin (E422) 49.850 mg/kg

ANALYTICAL COMPONENTS: Crude Protein 25.50% Crude Fat: 0.1% Cellulose: 2.78% Crude Ash: 1.1% Moisture: 73%

RECOMMENDED APPLICATION: Felaxa Lysine contains lysine which is an amino acid that strengthens the immune system, provides collagen formation, repairs tissue, supports antibody and white blood cell formation. It is used for supporting the treatment of upper respiratory tract diseases, conjunctivitis, herpes virus and strengthening the immune system in cats. It also contains malt and vegetable oils to facilitate the natural removal of ingested hairballs and other indigestible components. It doesn't contain artificial dyes or flavors.

USAGE AND DOSAGE: It can be given directly from the tube into the mouth or placed on the paw, or it can be added to the food. For adult cats: 2 cm/day
For kittens: 1 cm/day

PACKAGING: 80 g SHELF LIFE: 2 years







VITAMIN-MINERAL FOR RUMINANTS 1 kg - 5 kg - 20 kg

It is used by adding 1 kg to 100 kg feed to complete the vitamin and mineral deficiencies of lamb, sheep, kid, goat, calf, calve, cattle and bull.

USAGE AND DOSAGE: Viomix is added 1 kg to 100 kg feed. It should be mixed with a small portion of the feed; then it should be added slowly by mixing with the remaining feed. It should be ensured that the products is completely dispersed in the feed.



| Active materials | Level in Premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A | 250.000 | IU/kg |
| Vitamin D ₃ | 50.000 | IU/kg |
| Vitamin E / α-Tocopherol Acetate | 50 | mg/kg |
| Vitamin B ₁ / Thiamine Monohydrate | 10 | mg/kg |
| Vitamin B ₃ / Niacin | 2.000 | mg/kg |
| Vitamin B ₇ / D-Biotin 2% | 4 | mg/kg |
| Trace minerals | | |
| Iron/Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| lodine/Calcium lodate 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| Manganese/Manganese Oxide 62 | 5.000 | mg/kg |
| Zinc/Zinc Oxide 72 | 5.000 | mg/kg |
| Selenium/Sodium Selenite 4.5% | 30 | mg/kg |
| Technological additives | | |
| Antioxidants | | |
| BHT/Butyl Hydroxytoluene | 100 | mg/kg |
| Sensory Additives | | |
| Aromatic Compounds | 2.000 | mg/kg |
| Carriers | | |
| Phosphorus / Di Calcium Phosphate 18% P - 25% Ca | 2.680 | mg/kg |
| Calcium | 3.722 | mg/kg |
| Calcium/ Calcium Carbonate 39% Ca | 353.163 | mg/kg |
| Sodium Bicarbonate/Sodium Bicarbonate | 5.000 | mg/kg |
| Sodium Chloride | 20.000 | mg/kg |

PACKAGING: 1 kg, 5 kg, 20 kg SHELF LIFE: 24 months.

120

LAMB-KID 5 g/day mixed with 0.5 kg feed

DAIRY COW

75 g/day

mixed with 7.5 kg feed

10 g/day mixed with 1 kg feed

50 g/day

mixed with

SHEEP-GOAT

CALF-CALVE BEEF CATTLE 25 g/day mixed with 2.5 kg feed

5 kg feed BULL

75 g/day mixed with 7,5 kg feed.

MINEMIX GOLD

78













VITAMIN - MINERAL - PROBIOTIC FOR RUMINANTS 1 kg - 5 kg - 20 kg - 25 kg

It is a premix that contains vitamins, minerals and propiotics, prepared for the completion of vitamin and mineral substance deficiencies of lambs, sheep, kids, goats, calfs, calves, cattle and bulls.

RECOMMENDED APPLICATION: Minemix Gold is added 1 kg to 100 kg feed for lamb, sheep, goat, calf, beef cattle, dairy cow and bull feeds. The product should first be mixed with a small portion of the feed, then the remaining feed should be added slowly by mixing. It should be ensured that the product is completely dispersed in the feed. If the product is to be added to the feed of each animal separately, the following amounts are given per animal.

:5 g/day 0,5 kg feed Lamb-Kid Sheep-Goat :10 g/day 1 kg feed Calf-Veal :25 g/ day 2,5 kg feed Beef Cattle :50 g/day 5 kg feed :75 g/ day 7,5 kg feed Dairy Cow



| Active materials | Level in Premix | Units |
|---|--------------------|--------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A / Retinyl acetate | 250.000 | IU/kg |
| Vitamin D3 / Cholecalsiferol | 50.000 | IU/kg |
| Vitamin E / α-Tocopherol acetate | 50 | mg/kg |
| Vitamin B ₁ /Tiyamin | 10 | mg/kg |
| Vitamin B ₃ / Niasin | 2.000 | mg/kg |
| Vitamin B ₇ / D-(+) Biotin | 4 | mg/kg |
| Trace minerals | | |
| Ferrous sulfate monohydrate %30 | 5.000 | mg/kg |
| Iodine/Calcium iodate %10 | 80 | mg/kg |
| Cobalt/Cobalt carbonate %5 | 20 | mg/kg |
| Copper/Copper sulfate pentahydrate %25 | 1.000 | mg/kg |
| Manganese/Manganese oxide %62 | 5.000 | mg/kg |
| Zinc/Zinc oxide %72 | 5.000 | mg/kg |
| Selenium/Sodium selenite %4,5 | 30 | mg/kg |
| Zootechnical feed additives / Intestinal flora regulators | | |
| Microorganisms | | |
| Saccharomyces cerevisae | 3x10 ¹¹ | cfu/kg |
| Preservatives | | |
| BHT | 100 | |
| Sensory Additives | | |
| Aromatic compounds | | |
| Aromatic mixture | 2.000 | mg/kg |
| Carriers | | |
| Di calcium phosphate | 15.000 | mg/kg |
| Sodium chloride | 20.000 | mg/kg |
| Sodium bicarbonate | 5.000 | mg/kg |
| Calcium carbonate | 900.000 | mg/kg |

PACKAGING: 1 kg, 5 kg, 20 kg, 25 kg

SHELF LIFE: 24 months

MINEMIX




VITAMIN - MINERAL - PROBIOTIC FOR RUMINANTS

1 kg - 5 kg - 20 kg









It is used by adding 2 kg to 100 kg of feed to meet the vitamin and mineral deficiencies of lambs, sheep, kids, goats, calves, cattle and bulls.

RECOMMENDED APPLICATION: The product should first be mixed with a small portion of the feed, then the remaining feed should be added slowly by mixing. It should be ensured that the product is completely dispersed in the feed. If the product is to be added to the feed of each animal separately, the following amounts are given per animal.

Lamb-Kid :10 g/day 0,5 kg feed Sheep-Goat :10 g/day 1 kg feed :50 g/day 2,5 kg feed Calf-Veal Beef Cattle :100 g/day 5 kg feed Dairy Cow :150 g/day 7,5 kg feed



| Active materials | Premiksteki Düzey | Birimler |
|---|--------------------|----------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A / Retinyl acetate | 125.000 | IU/kg |
| Vitamin D3 / Cholecalsiferol | 50.000 | IU/kg |
| Vitamin E / α-Tocopherol acetate | 50 | mg/kg |
| Vitamin B ₁ /Tiyamin | 10 | mg/kg |
| Vitamin B ₃ / Niasin | 1.000 | mg/kg |
| Vitamin B ₇ / D-(+) Biotin | 4 | mg/kg |
| Trace minerals | | |
| Ferrous sulfate monohydrate %30 | 2.000 | mg/kg |
| lodine/Calcium iodate %10 | 32 | mg/kg |
| Cobalt/Cobalt carbonate %5 | 8 | mg/kg |
| Copper/Copper sulfate pentahydrate %25 | 400 | mg/kg |
| Manganese/Manganese oxide %62 | 2.000 | mg/kg |
| Zinc/Zinc oxide %72 | 2.000 | mg/kg |
| Selenium/Sodium selenite %4,5 | 12 | mg/kg |
| Zootechnical feed additives / Intestinal flora regulators | | |
| Microorganisms | | |
| Saccharomyces cerevisae NCYC Sc 47/CNCM I-4407 | 2x10 ¹⁰ | cfu/kg |
| Preservatives | | |
| BHT | 100 | |
| Sensory Additives | | |
| Aromatic compounds | | |
| Aromatic mixture | 1.000 | mg/kg |
| Carriers | | |
| Di calcium phosphate | 2680 | mg/kg |
| Sodium chloride | 20.000 | mg/kg |
| Sodium bicarbonate | 5.000 | mg/kg |
| Calcium carbonate | 488.287 | mg/kg |

BİYO ANİSOL





Vitamins, provitamins and substances with chemically well-defined similar properties

ANALYTICAL COMPONENTS: Crude Protein 1%, Crude Fat 0%, Crude Cellulose 0%, Crude Ash 1.1%

RECOMMENDED APPLICATION: As a supportive it meets the vitamin and amino acid needs of all target animal species listed below. It is also used as a supplement in accelerating their development, increasing their productivity, increasing feed efficiency and strengthening

USAGE AND DOSAGE: It can be used by mixing drinking water of animals or by drinking directly. It should be ensured that animals

Animal species

Parrot and Pigeon Cat and Dog

Horse and Cattle

PACKAGING: 100 ml, 500 ml, 1000 r

SHELF LIFE: 24 months

Canary

Poultry

Budgerigar

Active material / Additive Name

Vitamin B₂ (Riboflavin-5 phosphate)

Active material / Additive Name

Active material / Additive Name

Amino acids Salts and analogs of amino acids:

consume the medicated water within 7-8 hours at least.

Vitamin B_s (Pantothenic acid)

Vitamin B₆ (Pyridoxine)

Vitamin B, (Tiamin)

Niasinamide

L- Tryptophan

DL-Methionine

Flavoring agents

Benzyl alcohol

Component List
Sodium chloride

Magnesium chloride

the immune system during illness.

Biotin

L-Lysine

L-Threonine

Choline chloride





SOLUTION VITAMIN - AMINO ACID 100 ml - 500 ml - 1000 ml

KALFOMIN

Code Level in premix

Code Level in premix

500 mg

500 mg

500 mg

1.000 mg

3 mg

2.000 mg

20.000 mg

1.500 mg

3.000 mg

8.000 mg

4.200 mg

10.000 mg

18.000 mg

170 mg

Level in premix

(3a831)

3a820

3a 826

3a841

3a315

3a890

3a880

3c440

3c301

3c321

3c410

2b02010

11.4.1

11.2.6

Dosage

1-2 drops

4-5 drops

2-3 ml/10 kg b.w

10-20 ml (hive)

10-20ml/100 kg b.w 1 lt/1000 lt drinking water





120

....

SOLUTION
LIQUID AMINO ACID AND MINERAL
1 lt - 20 lt

RECOMMENDED APPLICATION: KALFOMİN Solution is a balanced liquid mineral premix containing essential microelements and amino acids for the development and productivity of farm animals.

- · In poultry: It increases feed conversion ratio. It strengthens the egg shell, stops the mineral loss.
- · In Lambs and Calves: It eliminates micro element deficiency. It increases feed conversion ratio. It accelerates the development of young animals gain weight. It strengthens the immune system.
- In Adult Animals: It eliminates mineral deficiencies caused by nutritional and physiological reasons. It meets the increasing mineral need during pregnancy and lactation.

USAGE AND DOSAGE: Shake before use.

| Animal species | Dosage |
|--|-------------------------------------|
| Beef cattle | 20 ml /1 L water (for 100 kg bw) |
| Pregnant/ dairy cows | 30 ml / 1 L water (for 100 kg bw) |
| Calfs, pregnant and dairy sheeps and goats | 15 ml / 0,5 L water |
| Lambs and kids | 10 ml / 0,5 L water |
| Chicken and turkey | 1 L/ 200 L water (for 1000 chicken) |
| Chicks | 200-400 L for 1000 chicks |

| Active material | Level in Premix | Units |
|---|-----------------|-------|
| Nutritional Additives | | |
| Trace Elements | | |
| Iron/Iron III Chloride Anhydrous (34.38% Fe.) | 1.600 | mg/kg |
| Manganese/Manganese II Chloride 4 H2O (27.77% Mn) | 600 | mg/kg |
| Cobalt/Cobalt Sulfate .H2O (20% Co) | 250 | mg/kg |
| Zinc/Zinc II Chloride.H2O (47.97% Zn) | 650 | mg/kg |
| Copper/Copper II Chloride (36.5% Cu) | 250 | mg/kg |
| lodine/Potassium lodide (74% I) | 100 | mg/kg |
| Selenium/Sodium Selenite anhydrous 45% Se | 50 | mg/kg |
| Aminoacids | | |
| Lysine | 5.000 | mg/kg |
| DI-Methionine | 10.000 | mg/kg |
| Preservatives | | |
| Citric Acid | 10.000 | mg/kg |
| Phosphoric Acid | 13.000 | mg/kg |
| Carriers | | |
| Deionized water | 95.24 | % |

PACKAGING: 1 lt, 20 lt SHELF LIFE: 24 months







віуоміх



COMPLEMENTARY FEED FOR RUMINANT 5 kg - 20 kg

It is used by adding 1 kg to 100 kg feed to complete the vitamin and mineral substance deficiencies of lambs, sheep, kids, goats, calves, bulls.



| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A | 1.000.000 | mg/kg |
| Vitamin D ₃ | 100.000 | mg/kg |
| Vitamin E/α-Tocopherol Acetate | 500 | mg/kg |
| Vitamin B _/ /Thiamine Monohydrate | 100 | mg/kg |
| Vitamin B ₂ /Riboflavin | 100 | mg/kg |
| Vitamin B ₃ /Niacin | 1.000 | Mg/kg |
| Vitamin B _e /Pridoxine HCL | 100 | mg/kg |
| Vitamin B ₇ /d-Biotin 2% | 6 | mg/kg |
| Trace minerals | | |
| Iron /Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| lodine/Calcium lodate 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| Manganese/Manganese Oxide 62% | 5.000 | mg/kg |
| Zinc/Zinc Oxide 72% | 5.000 | mg/kg |
| Molybdenum/Sodium Molybdate 39% | 10 | mg/kg |
| Selenium/Sodium Selenite 4.5% | 30 | mg/kg |
| Technological additives | | |
| Antioxidants | | |
| BHT/Butyl Hydroxytoluene | 1000 | mg/kg |
| Sensory Additives | | |
| Aromatic Compounds | 1.000 | mg/kg |
| Carriers | | |
| Phosphorus/Di Calcium Phosphate 18% P - 25% Ca | 2.700 | mg/kg |
| Calcium | 3.750 | mg/kg |
| Calcium/Calcium Carbonate 39% Ca | 357.679 | mg/kg |
| Sodium Bicarbonate/Sodium bicarbonate | 5.000 | mg/kg |
| Sodium Chloride/Sodium Chloride | 20.000 | mg/kg |

PACKAGING: 5 kg, 20 kg SHELF LIFE: 2 years

120

BIYOSACC

LAMB-CALF SHEEP-GOAT

5 g/day mixed 10 g/day mixed

25 g/day mixed 50 g/day mixed

with 2,5 kg feed. with 5 kg feed.

50 g/day mixed 75 g/day mixed

with 5 kg feed. with 7,5 kg feed.

with 1 kg feed.

BEEF CATTLE

BULL

with kg feed.

CALF

DAIRY COW



VITAMIN, MINERAL, PROBIOTIC PREMIX 5 kg - 20 kg

BİYOSACC is a balanced feed supplement containing vitamins, minerals and probiotic (Saccharomyces cerevisiae) for large and small ruminants.

USAGE AND DOSAGE: 1 kg product is added to 100 kg of lamb, sheep, goat, calf, beef cattle, dairy cow and bull feeds. It should be mixed with a small portion of the feed; then it should be added slowly by mixing with the remaining feed. It should be ensured that the products is completely dispersed in the



| Active material | Level in Premix | Units |
|--|--------------------|--------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A | 1.000.000 | IU/kg |
| Vitamin D ₃ | 100.000 | IU/kg |
| Vitamin E/α-Tocopherol Acetate | 500 | mg/kg |
| Vitamin B ₁ /Thiamine Monohydrate | 100 | mg/kg |
| Vitamin B ₂ /Riboflavin | 100 | mg/kg |
| Vitamin B ₃ /Niacin | 1.000 | mg/kg |
| Vitamin B ₆ /Pridoxine HCL | 100 | mg/kg |
| Vitamin B ₇ /Biotin 2% | 6 | mg/kg |
| Trace minerals | | |
| Iron/Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| lodine/Calcium lodate 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| Manganese/Manganese Oxide 62% | 5.000 | mg/kg |
| Zinc/Zinc Oxide 72% | 5.000 | mg/kg |
| Selenium/Sodium Selenite 45% | 30 | mg/kg |
| Technological and Zootechnical Additives | | |
| Microorganisms | | |
| Saccharomyces Cereviase NCYC Sc 47 | 4x10 ¹⁰ | cfu/kg |
| Antioxidants | | |
| BHT/Butyl Hydroxytoluene | 1000 | mg/kg |
| Sensory additives | | |
| Aromatic compounds | 1.000 | mg/kg |
| Carriers | | |
| Phosphorus/Di Calcium Phosphate 18% P - 25% Ca | 2.700 | mg/kg |
| Calcium | 3.750 | mg/kg |
| Calcium/Calcium Carbonate 39% Ca | 356.119 | mg/kg |
| Sodium Bicarbonate/Sodium bicarbonate | 5.000 | mg/kg |
| Sodium Chloride/Sodium Chloride | 20.000 | mg/kg |

PACKAGING: 5 kg, 20 kg SHELF LIFE: 24 months

5 g/day to 0,5 kg feed

SHEEP - GOAT 10 g/day to 1 kg

BEEF CATTLE CALF 25 g/day to 5 kg feed 2,5 kg feed

DAIRY COW 50 g/day to 5 kg feed

75 g/day to 7,5 kg feed

VIOMIN-B



COMPLEMENTARY FEED FOR RUMINANT

1 kg - 5 kg - 25 kg

It is used lamb, sheep, calve, heifer, steer, cattle, bull and goat by adding 10 kg to 1 ton feed to meet LAMBS AND KIDS SHEEP - GOAT

| min and mineral deficiencies. | | |
|--|-----------------|-------|
| active material | Level in Premix | Units |
| eed additives | | |
| 'itamins | | |
| litamin A | 1.000.000 | mg/kg |
| fitamin D ₃ | 100.000 | mg/kg |
| 'itamin E/α-Tocopherol Acetate | 1.200 | mg/kg |
| itamin C/Ascorbic Acid | 100 | mg/kg |
| 'itamin B ₁ /Thiamine Monohydrate | 60 | mg/kg |
| fitamin B ₂ /Riboflavin | 225 | mg/kg |
| fitamin B ₃ /Niacinamide | 600 | mg/kg |
| fitamin B _s /Calcium D-pantetonate | 400 | mg/kg |
| 'itamin B _e /Pridoxine HCL | 10 | mg/kg |
| fitamin K ₃ /Menadione Sodium Bisulfite | 100 | mg/kg |
| race elements | | |
| ron/Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| odine/Calcium Iodate BMP 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate BMP 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| langanese/Manganese Oxide 62 | 5.000 | mg/kg |
| linc/Zinc Oxide 72 | 5.000 | mg/kg |
| elenium/Sodium Selenite BMP 4.5% | 30 | mg/kg |
| ensory Additives | | |
| romatic Compounds | | |
| aromatic mixtures | 2.000 | mg/kg |
| arriers | | |

PACKAGING: 1 kg, 5 kg, 25 kg SHELF LIFE: 24 months

Sodium Chloride

120

Calcium/Calcium Carbonate 39% Ca

Phosphorus/Di Calcium Phosphate 18% P - 25% Ca



39.000

54.200

185.250

250.000

mg/kg

mg/kg

mg/kg

VIOMIN-E

10 g/day

1 kg feed

BEEF CATTLE

50g/day

5 kg feed

BULL

75 g/day

7.5 kg feed.

5 g/day

0,5 kg feed

CALF

25 g/day

2,5 kg feed

DAIRY COW

50 g/day

5 kg feed



VITAMIN, MINERAL, AMINOACID PREMIX FOR RUMINANTS 25 kg

USAGE AND DOSAGE: It is a vitamin-mineral-amino acid premix for calves, cows, bulls, lambs, kids, sheep and goats at the rate of 1 kg per 100 kg of feed. It is used to meet the vitamin, mineral and amino acid needs of animals and to increase feed utilization.

Viomin E is added 1 kg to 100 kg feed. Viomin E should be mixed with a small portion of the feed first, then the remaining feed should be added slowly by stirring and it should be ensured that it is fully distributed into the feed.



| Active material | Level in Premix | Units |
|---|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin A | 2.000.000 | IU/kg |
| Vitamin D ₃ | 200.000 | IU/kg |
| Vitamin E / α-Tocopherol Acetate | 20.000 | mg/kg |
| Vitamin C / Ascorbic Acid | 4.000 | mg/kg |
| Vitamin B ₁ / Thiamine Monohydrate | 350 | mg/kg |
| Vitamin B ₂ / Riboflavin | 900 | mg/kg |
| Vitamin B ₃ / Niacin | 2.500 | mg/kg |
| Vitamin B _s / Calcium-d-Pantothenate | 1.000 | mg/kg |
| Vitamin B ₆ / Pyridoxine HCL | 90 | mg/kg |
| Vitamin B ₇ / D-Biotin 2% | 15 | mg/kg |
| Vitamin B₀/ Folic Acid | 350 | mg/kg |
| Vitamin B ₁₂ / Cyanocobalamin | 1,5 | mg/kg |
| Vitamin K ₃ / Menadione Sodium Bisulfite | 4.000 | mg/kg |
| Amino Acids | | |
| dl-Methionine | 5.000 | mg/kg |
| I-Lysine | 7.000 | mg/kg |
| Trace minerals | | |
| Iron/Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| Iodine/Calcium Iodate 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| Manganese/Manganese Oxide 62 | 5.000 | mg/kg |
| Zinc/Zinc Oxide 72 | 5.000 | mg/kg |
| Selenium/Sodium Selenite 4.5% | 30 | mg/kg |
| Technological additives | | |
| Antioxidants | | |
| Ethoxyquin | 314,4 | mg/kg |
| Propil Galat | 64,8 | mg/kg |
| Preservatives | | |
| Citric Acid | 222 | mg/kg |
| Binders and anti-caking agents | | |
| Colloidal Silica | 292,8 | mg/kg |
| Carriers | | |
| Phosphorus / Sodium Phosphate 12.89% P | 57,8 | mg/kg |
| Phosphorus/Dicalcium Phosphate 18% P-25% Ca | 39.000 | mg/kg |
| Calcium | 54.000 | mg/kg |
| Calcium/ Calcium Carbonate 39% Ca | 259.480 | mg/kg |
| Sodium Chloride | 30.000 | mg/kg |

PACKAGING: 25 kg SHELF LIFE: 24 months. LAMB-KID 5 g/day mixed with

25 g/day

50 g/day

5 kg feed

0.5 kg feed CALF-CALVE

BEEF CATTLE 25 g/day mixed with mixed with 2.5 kg feed

5 kg feed DAIRY COW BULL mixed with

75 g/day mixed with 5 kg feed

SHEEP-GOAT

10 g/day

mixed with

1 kg feed

20

VITHORSE











VITAMIN - MINERAL - AMINOACID 25 kg

USAGE AND DOSAGE: It is a premix prepared to be added to the feed of foals, mares, pregnant mares, horses and stallions at a rate of 10 kg ton.

Vithorse is added 1 kg to 100 kg feed. Vithorse should be mixed with a small portion of the feed first, then the remaining feed should be added slowly by stirring and it should be ensured that it is fully distributed into the feed.

| 3-6 month old foals | 10-15 g/day mixed with 1-1,5 kg feed. |
|--|---------------------------------------|
| Adult foals | 25 g/day mixed with 2.5 kg feed. |
| Pregnant and lactating mares and stallions | 50 g/day mixed with 5 kg feed. |
| Race horses | 75 g/day mixed with 7,5 kg feed. |
| | J. , |

| Active material | Level in Premix | Birimler |
|---|-----------------|----------|
| Nutritional Additives | | |
| Vitamins | | |
| Vitamin A | 2.000.000 | IU/kg |
| Vitamin D3 | 200.000 | IU/kg |
| Vitamin E / α-Tocopherol Acetate | 20.000 | mg/kg |
| Vitamin C / Ascorbic Acid | 4.000 | mg/kg |
| Vitamin B, / Thiamine Monohydrate | 350 | mg/kg |
| Vitamin B ₂ / Riboflavin | 900 | mg/kg |
| Vitamin B ₃ / Niacin | 2.500 | mg/kg |
| Vitamin B _s / Calcium-d-Pantothenate | 1.000 | mg/kg |
| Vitamin B ₆ / Pyridoxine HCL | 90 | mg/kg |
| Vitamin B ₇ / D-Biotin 2% | 15 | mg/kg |
| Vitamin B ₉ / Folic Acid | 350 | mg/kg |
| Vitamin B ₁₂ / Cyanocobalamin | 1,5 | mg/kg |
| Vitamin K ₃ / Menadione Sodium Bisulfite | 4.000 | mg/kg |
| Aminoacids | | |
| dl-Methionine | 5.000 | mg/kg |
| l-Lysine | 7.000 | mg/kg |
| Trace elements | | |
| Iron/Iron Sulfate Monohydrate 30% | 5.000 | mg/kg |
| lodine/Calcium Iodate 10% | 80 | mg/kg |
| Cobalt/Cobalt Carbonate 5% | 20 | mg/kg |
| Copper/Copper Sulphate Pentahydrate 25% | 1.000 | mg/kg |
| Manganese/Manganese Oxide 62 | 5.000 | mg/kg |
| Zinc/Zinc Oxide 72 | 5.000 | mg/kg |
| Molybdenum/Sodium Molybdate 39 | 10 | mg/kg |
| Selenium/Sodium Selenite 4.5% | 30 | mg/kg |
| Technological Contributions | | |
| Antioxidants | | |
| Ethoxyquin | 314,4 | mg/kg |
| Propil Galat | 64,8 | mg/kg |
| Preservatives | | |
| Citric acid | 222 | mg/kg |
| Binders and anti-caking agents | | |
| Colloidal Silica | 292,8 | mg/kg |
| Carriers | | |
| Phosphorus / Sodium Phosphate 12.89% P | 57,8 | mg/kg |
| Phosphorus/Dicalcium Phosphate 18% P-25% Ca | 39.000 | mg/kg |
| Calcium | 54.000 | mg/kg |
| Calcium/ Calcium Carbonate 39% Ca | 259.480 | mg/kg |
| Sodium Chloride | 30.000 | mg/kg |



BIYO EVIT-SE



VITAMIN - MINERAL COMPLEMENTARY FEED 100 g

RECOMMENDED APPLICATION: It is a complementary feed which eliminates the deficiency of Vitamin E and selenium minerals. It can be added to the milk or drinking water of lambs, calves, kids, calf and can be added drinking water of sheep, goats, cattle, chickes, chickens, turkeys.

USAGE AND DOSAGE:

- · For cage birds and pigeons, 4 g for per 1 liter of drinking water
- · For poultry 100 g for per 1 tin (25 kg) drinking water
- · For lambs, kids, calves, 4g is added to milk or drinking water for 20 kg body weight.
- · Sheep, goat, cattle: 10 g for 40-50 kg body weight to drinking water

These amounts can be repeated 4-5 times at 2-3 week intervals or as long as needed.

| Analytical components | |
|-----------------------|-------|
| Crude Protein | % 0,2 |
| Crude cellulose | % O |
| Oil (total) | % 2,1 |
| Crude Ash | % 0,8 |

| Active material | Le | vel in Premix | Units |
|-----------------------------|----|---------------|-------|
| Feed additives | | | |
| Vitamins | | | |
| Vitamin E/Vitamin E Acetate | | 30.000 | mg/kg |
| Trace minerals | | | |
| Sodium Selenite %45 | | 230 | mg/kg |
| Carriers | | | |
| Lactose/Lactose Monohydrate | | 970.000 | mg/kg |
| | | | |

PACKAGING: 100 g net SHELF LIFE: 24 months







BİYO AVİT-K











VITAMIN COMPLEMENTARY FEED

100 g - 1 kg

RECOMMENDED APPLICATION: It is a complementary feed that eliminates the deficiency of vitamins A and K3 that prepared to be added to the drinking water in lamb, calf, kid, calve and foal and to the milk or drinking water in poultry such as chicks, chickens and turkeys.

USAGE AND DOSAGE:

- $\cdot \text{For chicks, chickens, turkeys and other poultry animals: 100 g for 400 liters of drinking water or 20 kg for 20 kg for 20$
- \cdot 100 g for 200 liters of drinking water or 8 g for 20 liters of drinking water if they are required to adapt to the adverse conditions in the environment
- · For lambs, kids, calf and calves: Add 4 g per day for 10 lambs, kids, 4 g per day for 2 calves to 5-10 liters of milk or water.
- · 10 calves, foals: 8 g for 5-10 liters of water

| 0,2 |
|-----|
| 60 |
| 2,3 |
| 0,5 |
| |

| Active material | Level in | Units |
|--|------------|-------|
| Feed additives | Premix | |
| Vitamins | | |
| Vitamin A | 20.000.000 | IU/kg |
| Vitamin K_s /Menadione Sodium Bisulfite | 10.000 | mg/kg |
| Carriers | | |
| Lactose/Lactose Monohydrate | 950.000 | mg/kg |

PACKAGING: 100 g, 1 kg SHELF LIFE: 24 months





VIOSOL-K









MULTIVITAMIN POWDER FOR POULTRY 50 g - 100 g


RECOMMENDED APPLICATION: It is a vitamin premix for the daily vitamin needs of poultry. It is used by adding 2 g to 1 liter of drinking water and 2 kg to 1 ton of drinking water for the daily vitamin needs of chicks, chickens, turkeys, geese and ducks.

| Active mate | rial | Level in Premix | Units |
|-------------------------|----------------------------|-----------------|-------|
| Nutritional s | supplements | | |
| Vitamins | | | |
| Vitamin A | Retinol | 5.000.000 | IU/kg |
| Vitamin D ₃ | Cholecalciferol | 500.000 | IU/kg |
| Vitamin E | α-tocopherol acetate | 4.000 | mg/kg |
| Vitamin C | Askorbic acid | 10.000 | mg/kg |
| Vitamin B ₁ | Thiamine HCI | 2.000 | mg/kg |
| Vitamin B ₂ | Riboflavin-5-phosphate | 2.500 | mg/kg |
| Vitamin B ₆ | Pyridoxine HCl | 2.000 | mg/kg |
| Vitamin B ₁₂ | Cyanocobalamin | 5 | mg/kg |
| Vitamin B _s | Calcium-d-pantotenat | 5.000 | mg/kg |
| Vitamin B ₃ | Niacinamide | 10.000 | mg/kg |
| Vitamin K ₃ | Menadione sodium bisulfite | 2.000 | mg/kg |
| Vitamin B ₉ | Folic acid | 500 | mg/kg |
| Vitamin B ₇ | d-Biotin | 40 | mg/kg |
| Carriers | | | |
| Lactose | | 943.200 | mg/kg |

PACKAGING: 50 g, 100 g SHELF LIFE: 24 months







PRODIGEST



1,40







INTESTINAL FLORA REGULATOR PROBIOTIC AND
MINERAL PREMIX
190 g

RECOMMENDED APPLICATION: It is a intestinal flora regulating probiotic and mineral mixture that prepared to be added to the water of kids and calves, and to the feed of cattle, sheep and goats. 190 g Prodigest is given in 5 liters of water and mixed with 5 kg of food.

| Active material | Level in Premix | Units |
|--|--------------------|--------|
| Technological additives | | |
| Preservatives | | |
| Sodium propionate | 320.000 | mg/kg |
| Calcium propionate | 220.000 | mg/kg |
| Sensory Additives | | |
| Aromatic Compounds | | |
| Aromatic mixtures | 2.000 | mg/kg |
| Nutritional Additives | | |
| Trace Elements | | |
| Iron/Iron sulfate monohydrate (Fe 32.87%) | 580 | mg/kg |
| Cobalt/Cobalt sulfate heptahydrate (Co 34.07%) | 136 | mg/kg |
| Copper/Copper sulfate pentahydrate (Cu 25%) | 350 | mg/kg |
| Manganese/Manganese sulfate monohydrate (Mn 32.5%) | 400 | mg/kg |
| Zinc/Zinc sulfate monohydrate (Zn 36.44%) | 20 | mg/kg |
| Zootechnical additives | | |
| Intestinal flora regulators | | |
| Saccharomyces cerevisiae NCYC Sc 47 | 4X10 ¹⁰ | CFU/kg |
| Carriers | | |
| Sodium chloride | 103.000 | mg/kg |
| Sodyum karbonat | 350.000 | mg/kg |

PACKAGING: 190 g SHELF LIFE: 24 months





BİYO SEL-DE CARLSBAD



POWDER COMPLEMENTARY FEED250 g - 500 g

RECOMMENDED APPLICATION: It is a complementary feed for sheep, goats, cattle, horses, dogs and cats.

 Cattle and Horse
 : 75-100 g /day

 Sheep and Goat
 : 15-20 g / day

 Cat and dog
 : 5-10 g / day

| Analytical components | |
|-----------------------|---------|
| Calcium | %0,035 |
| Phosphorus | % O |
| odium | % 30,99 |
| Magnesium (1997) | % 0,006 |
| Crude ash | > % 10 |
| | |

| ctive material | Level in | Units |
|-------------------------|----------|-------|
| ineraller | Premix | |
| 4.6 Sodium Sulfate | 440.000 | mg/kg |
| 5.2. Potassium Sulfate | 20.000 | mg/kg |
| 4.2. Sodium Bicarbonate | 360.000 | mg/kg |
| 4.1. Sodium chloride | 180.000 | mg/kg |
| | | |

PACKAGING: 250 g, 500 g **SHELF LIFE:** 24 months









BİYO MAGNESİE CALCINEI









POWDER COMPLEMENTARY FEED 250g - 500 g

It is a complementary feed used for magnesium supplementation in sheep, goats, cattle and horses by adding to water. It supports the regulation and stabilization of rumen ph.

USAGE AND DOSAGE:

Veal and Foal : 20-25 g/animal/day Cattle and Horse : 50-75 g/animal/daY Sheep and Goats : 15-20 g /animal/day

| Analytical components | |
|-----------------------|--------|
| Calcium | %0,77 |
| Phosphorus | %0 |
| Sodium | %0 |
| Magnesium | %56,72 |
| Crude Ash | >15 |

| Level in Premix | |
|-----------------|-------|
| | Units |
| 1.000.000 | mg/kg |
| | |

PACKAGING: 250 g, 500 g SHELF LIFE: 24 months







BİYO MAGNEZYUM SÜLFAT





120

COMPLEMENTARY FEED 250 g - 500 g

RECOMMENDED APPLICATION: It is used for magnesium supplementation in sheep, goats, cattle and horses.

Sheep and Goat : 7.5-15 g/day : 60-120 g/day Calf and Foal :10-20 g/day : 30-60 g/day : 2-6 g/day : 1-3 g/day

PACKAGING: 250 g , 500 g SHELF LIFE: 24 months

| Analytical components | |
|-----------------------|-------|
| Calcium | %0 |
| Phosphorus | %0 |
| Sodium | %0 |
| Magnesium | %10,2 |
| Sulfur | %15 |
| Crude Ash | >%15 |

| Active material Minerals | Level in Premix | Units |
|--------------------------|-----------------|-------|
| Magnesium Sulfate / | | |
| Magnesium Sulfate | 1.000.000 | mg/kg |
| Heptahydrate | | |

























COMPLEMENTARY FEED 500 g

BİYOCAL Solution is a complementary liquid mineral feed used in calcium and magnesium deficiency of lambs, kids, sheep, goats, calves and cattle.

USAGE AND DOSAGE: It is drunk in the following amounts to meet the increased calcium requirement of pregnant cows during pregnancy and after calving:

 $1\,bottle\,approximately\,24\,hours\,before\,birth/1\,bottle\,immediately\,after\,birth\,1\,bottle\,12\,hours\,after\,birth/1\,bottle\,24\,hours\,after\,birth.$ It is used in calcium and magnesium deficiency of lamb, kid, sheep, goat, calf, calf and cattle.

1/4 bottle for 25 kg body weight, 3/4 bottle for 75 kg body weight

½ bottle for 50 kg body weight, 1 bottle for 100 kg body weight.

It is recommended to drink once a monthdepend on the condition of the animal.

PACKAGING: 500 g SHELF LIFE: 24 months



| Analytical components | |
|---------------------------|---------|
| Calcium | %10,5 |
| Phosphorus | % O |
| Sodium | % 0,011 |
| Magnesium | % 25 |
| Crude ash (550 C°3 hours) | % 1 |

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Calcium Chloride/Calcium Chloride 2H ₂ O | 313.382 | mg/kg |
| Magnesium Chloride/Magnesium Chloride 6 H ₂ O | 66.770 | mg/kg |
| Sodium Chloride/Sodium Chloride | 240 | mg/kg |
| Mono Propylene Glycol/Mono Propylene Glycol | 326.900 | mg/kg |
| | | |





BISMOLIN





RECOMMENDED APPLICATION: Bismolin Powder is a complementary feed that ensures healthy digestive functions of small and large ruminants and horses.

USAGE AND DOSAGE:

Calve-Calf-Foal Cow-Bull-Horse :50 g Lamb-Kid :5g :10 g Sheep-Goat

It is taken orally by mixing with water or milk at a ratio of 5 times.

PACKAGING: 50 g SHELF LIFE: 24 months

| Active material | Level in Premix | Units |
|---|-----------------|-------|
| Nutritional Additives | | |
| Vitamins | | |
| Vitamin B ₁ /Thiamine Monohydrate | 60 | mg/kg |
| Vitamin B ₂ /Riboflavin | 225 | mg/kg |
| Vitamin B ₃ /Niacin | 600 | mg/kg |
| Vitamin B _s /Calcium-D-Panthotenate | 400 | mg/kg |
| Vitamin B ₆ /Pridoxine HCL | 10 | mg/kg |
| Vitamin K ₃ /Menadione Sodium Bisulfite | 100 | mg/kg |
| Technological Additives | | |
| Preservatives | | |
| Sodium Methyl 4-Hydroxybenzoate | 2000 | mg/kg |
| Binders, Anti-caking Agents | | |
| Kaolin | 400.000 | mg/kg |
| Bentonite | 20.000 | mg/kg |
| Sorbitol Powder | 70.000 | mg/kg |
| Zootechnical Additives | | |
| Intestinal Flora Regulators | | |
| Enterococcus Faecium NCIMB 10415 2x10 ¹⁰ Cfu/g | 500 | mg/kg |
| Saccharomyces Cervisiae NCYC Sc 47 | 10.000 | mg/kg |
| Fillers | | |
| Pectin | 40.000 | mg/kg |
| Magnesium Stearate | 2.000 | mg/kg |
| Lactose/Lactose Monohydrate | 454.105 | mg/kg |

| Analytical Components | |
|------------------------------|-------|
| Crude Protein | % 0,5 |
| Crude Ash | % 34 |
| Crude Cellulose | % 2 |
| Crude Oil (total) | % 1,1 |

Component List

Pectin, Magnesium stearate, Lactose, Kaolin, Bentonit





120





ARISOL





MULTIVITAMIN FOR HONEYBEES
100 g

RECOMMENDED APPLICATION: It helps to obtain more efficiency from the hives and to adapt to the environment, environment and climate changes during the transportation of the hives. It is used as a vitamin supplement to maintain the number of bees in the hive in winter. For honey bees, 2 g is added to 1 liter of water. In normal cases such as obtaining more efficiency from the hives; 10 g is added to 5 liters of water. 10 g is added to 5 liters for "syrup" which is prepared with sugar, honey and water in spring and autumn.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Nutritional supplements | | |
| Vitamins | | |
| Vitamin A/ Retinyl Acetate | 1.800.000 | IU/kg |
| Vitamin D ₃ | 270.000 | IU/kg |
| Vitamin E/ α-tocopherol acetate | 280 | mg/kg |
| Vitamin C/Ascorbic acid | 3.000 | mg/kg |
| Vitamin B ₁ /Thiamine HCl | 800 | mg/kg |
| Vitamin B ₂ /Riboflavin-5-phosphate | 1.250 | mg/kg |
| Vitamin B ₃ /Niacinamide | 7.500 | mg/kg |
| Vitamin B _s /Calcium-d-pantothenate | 1.200 | mg/kg |
| Vitamin B ₆ /Pyridoxine HCL | 200 | mg/kg |
| Vitamin B ₇ /d-Biotin 2% | 10 | mg/kg |
| Vitamin B _q /Folic acid | 250 | mg/kg |
| Vitamin B ₁₂ /Siocobalamin | 1 | mg/kg |
| Vitamin K ₃ /Menadione sodium bisulfite | 1.500 | mg/kg |
| Carriers | | |
| Saccharose | 970.400 | mg/kg |

PACKAGING: 100 g SHELF LIFE: 24 months







BITTER BEAK



ANTI CANNIBAL SPRAY 100 ml - 250 ml

COMPOSITION: Each 100 ml spray contains 0.017 g Denatonium Benzoate.

RECOMMENDED APPLICATION: Although feather plucking in birds develops as a result of nutritional disorders and hormonal effects, or as a reaction to infestations caused by lice (mite), other parasites, it is often observed as a reaction to sedentary living conditions. Due to its formulation, it has a very bitter taste and is used to prevent feather plucking seen in parrots, other cage, perch and large cage birds. It prevents birds from biting their wounds.

USAGE AND DOSAGE: It is gently applied to the bird's affected feather area, around the wound, and to the dressing area. A once-daily application usually prevents the plucking problem. But if the problem persists, it is repeated as long as necessary. Spray it on your bird's feathers away from feeders and drinkers, as it tastes very bitter.

PACKAGING: 100 ml, 250 ml SHELF LIFE: 3 years





AVISENCID



SKIN CARE SOLUTION 10 cc - 20 cc - 30 cc


COMPOSITION: It is a solution containing Tea Tree Oil, Cinnamon Oil, Neem Oil, and Lavender Oil.

RECOMMENDED APPLICATION: It is a skin care product which used for cage birds such as canaries, budgies, parrots and pigeons. With the moisturizing effect of the essential oils it contains, it helps maintaining pH/moisture balance of the skin. It provides a soft, bright and natural vitality without irritating the skin. It is suitable for use on skin that has allergic reactions to flea, tick and lice bites. It relieves skin irritations caused by fleas, ticks and lice. It helps the main treatment in eliminating crusts, warts and similar formations that occur as a result of some skin diseases. Without affecting the normal flora of the bird, it removes bad odor and makes it smell good.

USAGE AND DOSAGE: It is applied topically on the skin. Avisencid 1 drop per 60 g weight, dropped between the shoulders on the back of the neck of the bird.

PACKAGING: 10 cc, 20 cc ve 30 cc

SHELF LIFE: 2 years







BIYO DERMACURE HERBIO AVISPREY









PUMP LIQUID SPRAY SKIN CARE CREAM 50 ml - 100 ml - 150 ml - 200 ml

COMPOSITION: 1 g of HERBIO Avispray is a colorless solution containing 40 mg of Palmarosa oil.

RECOMMENDED APPLICATION: Biyo Dermacure HERBIO Avispray contains natural Palmarosa oil derived from the Cymbopogon martinii plant. Palmarosa oil, which is used in alternative treatment in the east, especially in India, provides a healthy and shiny skin of cage birds and pigeons with its moisturizing feature. It maintains the natural oil balance of the skin. Bio Dermacure HERBIO Avispray also helps to treat wounds and bruises caused by self-pecking and plucking feathers (Cannibalism) of cage birds. It relieves skin irritations caused by fleas, ticks and lice.

USAGE AND DOSAGE: It is sprayed on the skin from a distance of 15 cm to cage birds and pigeons. It can be repeated after 15 days if necessary.

PACKAGING: 50 ml, 100 ml, 150 ml, 200 ml SHELF LIFE: 2 years

















SOLUTION 30 cc Avi-ast Solution meets the vitamin C needs of cage birds and pigeons such as budgerigars, canaries, parrots and pigeons.

USAGE AND DOSAGE:

For a bird, 15 drops are added to a medium-sized drinker (100 cc). In pigeons: 30 drops are added to a medium-sized drinker (100 cc). The water of the birds should changed once every two days.

| Active material | Level in Premix | Units |
|---|-----------------|-------|
| Nutritional supplements | | |
| Vitamins | | |
| Vitamin C/Ascorbic Acid | 7.500 | mg/kg |
| Natural Additives and Feed Ingredients | | |
| Propolis | 12.000 | mg/kg |
| Eucalyptus Oil | 12.000 | mg/kg |
| Preservatives | | |
| BHT/Butylhydroxy Toluene | 1.000 | mg/kg |
| Carriers | | |
| Glycerin | 20.000 | mg/kg |
| Deionized Water | 90 | % |
| | | |



PACKAGING: 30 cc SHELF LIFE: 24 months







AVI-DIAR

120

SOLUTION PREMIX 30 cc

Avi-diar Solution is an aromatic feed additive which uses for to meet the iron and zinc needs of cage birds and pigeons such as budgerigars, canaries, parrots.

| Active material | Level in Premix | Units |
|-------------------------------------|-----------------|-------|
| Trace elements | | |
| Ferrous/iron sulfate monohydrate | 2 | mg/kg |
| Zinc/ Zinc Sulfate Heptahydrate | 1,4 | mg/kg |
| Feed materials and Natural Products | | |
| Propolis | 20.000 | mg/kg |
| Pectin | 2.000 | mg/kg |
| Carriers | | |
| Glycerin | 20.000 | mg/kg |
| Deionized water | 96 | % |
| | | |

PACKAGING: 30 cc SHELF LIFE: 24 months







AVISOL-A



VITAMIN A - VITAMIN D3 - VITAMIN E VITAMIN C SOLUTION 20 cc

RECOMMENDED APPLICATION: AVISOL-A is used for the vitamin needs of ornamental and cage birds such as pigeons, budgies,

parrots and canaries, and positively affects their development.

For a bird, 2.5 cc is added to a medium size drinker (100 cc).

During the breeding and spawning periods of female and male birds;

For a bird, 5 cc is added to a medium sized drinker (100 cc).

At the time of bird moulting; For a bird, 5 cc is added to a medium sized drinker (100 cc).

The water of the birds should change every day.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Vitamins | | |
| Vitamin A | 400.000 | IU/kg |
| Vitamin D ₃ | 40.000 | IU/kg |
| Vitamin E/α-Tocopherol Acetate | 3.000 | mg/kg |
| Antioxidants | | |
| Vitamin C/Ascorbic Acid | 15.000 | mg/kg |
| BHT/Butyl Hydroxytoluene | 500 | mg/kg |
| Emulsifiers | | |
| Glyceryl polyethylene glycol ricinoleate | 50.000 | mg/kg |
| Carriers | | |
| Glycerin | 20.000 | mg/kg |
| Deionized water | 90,52 | % |
| | | |

PACKAGING: 20 cc SHELF LIFE: 24 months

. +







AVISOL-B

AVISOL - B
B-KOMPLEKS









VITAMIN SOLUTION FOR BIRDS 30 cc

AVISOL - B

RECOMMENDED APPLICATION: AVISOL-B is used for the vitamin needs of ornamental and cage birds such as pigeons, budgies, parrots and canaries, and positively affects their development.

For pigeon and big parrot; Add 5 cc to the medium size drinker (100 cc).

For other birds; Add 2.5 cc to a medium size drinker (100 cc).

The water of the birds should change every day.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin B ₁ /Thyamine Monohydrate | 1.000 | mg/kg |
| Vitamin B ₂ /Riboflavin | 1.000 | mg/kg |
| Vitamin B ₃ /Niacinemide | 5.500 | mg/kg |
| Vitamin B _s /d-Panthenol | 1.400 | mg/kg |
| Vitamin B ₆ /Pridoxine HCL | 1.000 | mg/kg |
| Vitamin B ₇ /d-Biotin 2% | 5,83 | mg/kg |
| Vitamin B ₁₂ /Cyanocobalamin | 0,52 | mg/kg |
| Stabilizers and Thickeners | | |
| Sorbitol | 105.000 | mg/kg |
| Glyceryl polyethylene glycol ricinoleate | 5.200 | mg/kg |
| Carriers | | |
| Propylene Glycol | 210.000 | mg/kg |
| Deionized Water | 70 | % |
| | | |

PACKAGING: 30 cc SHELF LIFE: 24 months







AVISOL-PLUS



VITAMIN SOLUTION FOR BIRDS

20 cc


RECOMMENDED APPLICATION: AVİSOL-Plus is used for the vitamin needs of ornamental and cage birds such as pigeons, budgies, parrots and canaries, and positively affects their development. It makes the colors look vivid and bright.

For a bird, 2.5 cc is added to a medium size drinker (100 cc).

During the breeding and spawning periods of female and male birds; For a bird, 5 cc is added to a medium sized drinker (100 cc). At the time of bird moulting; For a bird, 5 cc is added to a medium sized drinker (100 cc).

The water of the birds should change every day.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Vitamins | | |
| Vitamin A | 1.000.000 | IU/kg |
| Vitamin E/DL-α Tocopherol Acetate | 3.100 | mg/kg |
| Vitamin C/Ascorbic Acid | 20 | mg/kg |
| Vitamin B ₁ /Thiamine Mononitrate | 600 | mg/kg |
| Vitamin B ₂ /Riboflavin | 1.000 | mg/kg |
| Vitamin B ₃ /Niacinamide | 10.000 | mg/kg |
| Vitamin B _s /D-Panthenol | 2.500 | mg/kg |
| Vitamin B ₆ /Pridoxine HCL | 1.000 | mg/kg |
| Vitamin B ₇ /d-Biotin 2% | 25 | mg/kg |
| Vitamin B ₁₂ Cyanocobalamin | 4 | mg/kg |
| Emulsifiers | | |
| Glycerylpolyethylene glycol ricinoleate | 150.000 | mg/kg |
| Antioxidants | | |
| BHT/Butil Hidroksi Toluen | 1.000 | mg/kg |
| Carriers | | |
| Glycerin | 100.000 | mg/kg |
| Deionized water | 72,58 | % |



120

. +







AVISOL-SE





120

MINERAL SOLUTION FOR BIRDS 30 cc

RECOMMENDED APPLICATION: Avisol-Se is a feed additive that eliminates the deficiency of Vitamin E, Vitamin H (biotin) and selenium. It should be added to the drinking water of cage birds such as budgerigars, canaries, parrots and pigeons. Vitamin E is an antioxidant that ensures the health of the reproductive, muscular, circulatory, nervous and immune systems of animals. Giving vitamin E and selenium together as a supplement helps animals to adapt to the adverse conditions in their environment by increasing their body defense. Avisol-Se helps the offspring to grow and develop rapidly during laying and hatching of chicks.

In addition, because it contains Vitamin H (biotin), it prevents breakage in the claws and moulting.

USAGE AND DOSAGE: Add 1 cc (20 drops) to 100 cc drinker of cage birds such as budgerigars, canaries, parrots, and 1 cc (20 drops) for each pigeon. Should start 3-4 weeks before the breeding season and use it until laying and incubation periods. When spawning starts, the recommended dosage can be increased up to 2 times. The water of the birds should change every day.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Feed additives | | |
| Vitamins | | |
| Vitamin E/α Tocopherol Acetate Oily | 100.000 | mg/kg |
| Vitamin B ₇ /d-biotin 2% | 4.000 | mg/kg |
| Trace Elements | | |
| Sodium Selenite (selenium 45%) | 1,125 | mg/kg |
| Technological additives | | |
| Emulsifiers | | |
| Glyceryl Polyethylene Glycol Ricinoleate | 150.000 | mg/kg |
| Antioxidants | | |
| BHT/Butyl Hydroxy Toluene | 1.000 | mg/kg |
| Carriers | | |
| Glycerin | 100.000 | mg/kg |
| Deionized Water | 73,50 | % |
| | | |

PACKAGING: 30 cc SHELF LIFE: 24 months





AVIMINERAL



MINERAL SOLUTION FOR BIRDS 30 cc

RECOMMENDED APPLICATION: It meets the daily mineral needs of cage birds such as budgerigars, canaries, parrots and pigeons. In budgerigars, canaries, parrots, 30 drops (1.5 cc) are added to the amount of water that 1 bird can consume in a day (about 1/4 of a medium-sized drinker 25 cc) once a week. In pigeons; 100 drops (5cc) are added once a week to the water (50cc) that a pigeon can consume in 1 day.

| Active material | Level in Premix | Units |
|--|-----------------|-------|
| Trace minerals | | |
| ron/Iron Sulphate (Fe 32.87%) | 2 | mg/kg |
| odine/Potassium Iodide | 1 | mg/kg |
| Cobalt/Cobalt Sulfate (Co 34.07%) | 0,2 | mg/kg |
| Copper/Copper Sulphate (Cu 25%) | 0,2 | mg/kg |
| Manganese/Manganese Sulfate (Mn 32.5%) | 2 | mg/kg |
| Zinc/Zinc Sulphate (Zn 36.44%) | 1,4 | mg/kg |
| Selenium/Sodium Selenite 45 | 0,1 | mg/kg |
| Minerals | | |
| Magnesium /Magnesium Chloride | 0,010 | mg/kg |
| Calcium/Calcium Chloride | 0,6 | mg/kg |
| Sodium/Sodium Chloride | 0,08 | mg/kg |
| Sensory Additives | | |
| Colorants | | |
| Canthaxanthin | 50 | mg/kg |
| Carriers | | |
| Deionized water | 99,94 | % |

PACKAGING: 30 cc SHELF LIFE: 2 years

120







AVI-VIT



MULTIVITAMIN POWDER FOR BIRDS 50 g **RECOMMENDED APPLICATION:** It is a vitamin premix powder which prepared to meet the daily vitamin needs of cage birds and pigeons such as budgerigars, canaries, parrots and positively affect their development. 1 g is added to the medium size drinker (100 ml) of cage birds such as budgerigars, canaries and parrots. Add 2 g to 250 ml drinking water of pigeons.

WARNINGS: It should not be used other premixes containing Vitamin D2 and Vitamin D2 simultaneously.

| ctive Substance | Level in Premix | Units |
|--|-----------------|-------|
| lutritional Additives | | |
| 'itamins | | |
| itamin A | 4.000.000 | IU/kg |
| itamin D ₃ | 200.000 | IU/kg |
| itamin E/ Vitamin E Acetate | 4.000 | mg/kg |
| itamin C / Ascorbic Acid | 10.000 | mg/kg |
| itamin B ₁ /Thiamine Mononitrate | 2.000 | mg/kg |
| itamin B ₂ / Riboflavin | 2.500 | mg/kg |
| itamin B ₃ /Niacinamide | 10.000 | mg/kg |
| itamin B _s /Calcium D-Pantetonate | 3.000 | mg/kg |
| itamin B ₆ /Pridoxine HCL | 2.000 | mg/kg |
| itamin B ₇ / d-Biotin 2% | 40 | mg/kg |
| itamin B ₉ / Folic Acid | 500 | mg/kg |
| itamin B ₁₂ / Cyanocobalamin | 5 | mg/kg |
| itamin K ₃ / Menadione Sodium Bisulfite | 1.000 | mg/kg |
| arriers | | |
| actose/Lactose Monohydrate | 949.300 | mg/kg |
| | | |

PACKAGING: 50 g SHELF LIFE: 24 months







BİYO-AQUACURE CONTRABLUE



SOLUTION - AQUARIUM WATER REGULATOR 50 ml - 100 ml - 1000 ml

COMPOSITION: CONTRABLUE Solution contains Methylene Blue. It is a dark blue solution.

RECOMMENDED APPLICATION: CONTRABLUE Solution ensures that the aquarium stays clean. It prevents transmission of the disease agents to healthy fish. CONTRABLUE Solution is also oxygen carrier and transforms methemoglobin in the fish blood into normal oxygen. Therefore, it prevents poisoning of the fish by nitrite and/or cyanide probable to emerge in the aquarium.

USAGE AND DOSAGE:

- · Before the application, remove the active carbon filter.
- On the first day, add 5 ml CONTRABLUE solution per 20 liters of aquarium water. This provides the 3 ppm concentration required for cleaning of the aquarium water.
- On days 2 and 3, continue to add 2.5 ml CONTRABLUE Solution into the aquarium (do not attach the active carbon filter during this process).
- $\cdot \ \, \text{At the end of the day 3, refresh the aquarium water by discharging half of it, and attach the active carbon filter back.}$

PACKAGING: 50 ml, 100 ml, 1000 ml.

SHELF LIFE: 2 years





BİYO-AQUACURE CONTRA



SOLUTION - AQUARIUM WATER REGULATOR

30 ml - 50 ml - 100 ml - 1000 ml

COMPOSITION: CONTRA Solution contains 9-Amino-acridine HCl, Crystal violet, and Methylene Blue. It is a blue solution.

RECOMMENDED APPLICATION: CONTRA Solution ensures that the aquarium stays clean. It prevents transmission of the disease agents to healthy fish. CONTRA Solution is also oxygen carrier and transforms methemoglobin in the fish blood into normal oxygen. Therefore, it prevents poisoning of the fish by nitrite and/or cyanide probable to emerge in the aquarium.

USAGE AND DOSAGE:

- Add 10 cc CONTRA solution per 20 liters of aquarium water. This provides the concentration required for cleaning of the aquarium water.
- Repeat the application 3-4 times using the same amount every other day.
- After each application, keep the filter off for 3-4 hours to prevent discharge of the CONTRA solution from the aquarium water;
 you can turn on the filter again after this period.
- · At the end of an approximately I week application, replace the dirty filter material (sponge or active carbon) with a new one.

PACKAGING: 30 ml, 50 ml, 100 ml, 1000 ml .

SHELF LIFE: 2 years





BİYO-AQUACURE NİTROFUR



SOLUTION - AQUARIUM WATER REGULATOR

50 ml - 100 ml - 1000 ml

COMPOSITION: NITROFUR Solution is a yellow aquarium water regulator that contains Isopropyl alcohol, Carboxymethyl Cellulose, 5-nit-ro-2-furaldehyde semicarbazone, and Crospovidone.

RECOMMENDED APPLICATION: NİTROFUR Solution regulates the aquarium water and ensures that it stays clean by ensuring that the parasitic and fungal infections cannot propagate in the aquarium water.

USAGE AND DOSAGE:

- Add 5 ml NITROFUR solution per 20 liters of aquarium water. This provides the concentration required for cleaning of the aquarium water.
- Repeat the application for 3-4 days.
- After each application, keep the filter off for 3-4 hours to prevent discharge of the NITROFUR solution from the aquarium water; you can turn on the filter again after this period.
- · At the end of the application, replace the dirty filter material (sponge or active carbon) with a new one.
- $\,\cdot\,\,$ Repeat the application each time you refresh the aquarium water.

Nitrofur is an aquarium water regulator. It is not used in the pools and fish farms where fish is raised as food.

PACKAGING: 50 ml, 100 ml, 1000 ml.

SHELF LIFE: 2 years







REGIONAL CONTACT NUMBERS

Sales Manager : 0533 207 4882





SINCE 1955



BİYOTEKNİK A.Ş.

Head Office: Kocatepe Mah. Cumhuriyet Cad. Inkilap Apt. No: 15 D: 2 P.C.: 34437 Beyoglu - Istanbul / TÜRKİYE **Phone:** +90 0212 361 55 13

BT1 Factory: Kirklareli Organize Sanayi Bölgesi 5. Cad. No: 6/2 Kizilcikdere Koyu - Kırklareli / TÜRKİYE BT2 Factory: Kirklareli Organize Sanayi Bölgesi 5. Cad. No: 9 Kizilcikdere Koyu - Kırklareli / TÜRKİYE

www.biyoteknik.com

REGIONAL CONTACT NUMBERS

Sales Manager: 0533 207 4882 Aegean Region: 0537 397 5647

Southeastern Anatolia Region: 0533 202 3242

Black Sea Region: 0533 202 3240 Mediterranean Region: 0541 202 3249 Central Anatolia Region 0533 202 3242













